

中国昆虫病原线虫研究和产业化进展*

颜 珣^{1**} 郭文秀² 田成丽^{3,4} 李星月⁵ 陈书龙⁶ 金永玲⁷
李丽英⁸ 唐 睿⁸ 曹 莉^{8***} 韩日畴^{8***}

(1. 仲恺农业工程学院农业与生物学院, 农业农村部华南果蔬绿色防控重点实验室, 广州 510225; 2. 山东省农业科学院植物保护研究所, 山东省农业有害生物绿色防控重点实验室, 济南 250100; 3. 吉林省农业科学院(中国农业科技东北创新中心)农业资源与环境研究所, 长春 130033; 4. 辽宁大学生命科学院, 沈阳 110136; 5. 四川省农业科学院植物保护研究所, 农业农村部西南作物有害生物综合治理重点实验室, 成都 610066; 6. 河北省农林科学院植物保护研究所, 保定 071000; 7. 黑龙江八一农垦大学, 大庆 163319; 8. 广东省科学院动物研究所, 广东省动物保护与资源利用重点实验室, 广东省野生动物保护与利用公共实验室, 广州 510260)

摘 要 昆虫病原线虫(Entomopathogenic nematodes)作为国际上已经商业化应用、对土栖性和钻蛀性害虫特效的昆虫寄生性天敌,自20世纪60年代被陆续引入我国后,在农林和卫生等重要害虫绿色治理中发挥了重要作用。全球分离和鉴定了150多种昆虫病原线虫,我国发现的斯氏线虫*Steinernema*有27种,异小杆线虫*Heterorhabditis*6种,其中有19个斯氏线虫新种和2个异小杆线虫新种。对昆虫病原线虫共生细菌的研究,明确了共生细菌与昆虫病原线虫的共生专化性,并从共生细菌中鉴定了多种毒素和次生代谢物,具有杀虫、抗菌、杀线虫等生物活性,应用前景广阔。昆虫病原线虫田间应用的效果受环境因素的影响,通过对昆虫病原线虫环境耐性及机理的研究,筛选了适应环境压力的昆虫病原线虫品种和品系,应用于不同地区防控特定害虫。在昆虫病原线虫培养、剂型、保存和施用技术等方面开展的研发及技术创新工作,使昆虫病原线虫在我国实现了产业化。目前昆虫病原线虫已在我国推广应用,在防控桃小食心虫*Carposina nipponensis*、蛴螬、韭菜迟眼蕈蚊*Bradysia odoriphaga*幼虫等害虫上取得了良好的效果。本文从昆虫病原线虫鉴定和分类、昆虫病原线虫产业化、昆虫病原线虫共生细菌、昆虫病原线虫与共生细菌专化性、昆虫病原线虫应用、昆虫病原线虫环境耐受性、昆虫病原线虫与外界因子(包含生物与非生物因子)等领域全面综述我国昆虫病原线虫的研发进展。

关键词 昆虫病原线虫; 共生细菌; 鉴定分类; 产业化培养; 剂型; 田间应用

Research and commercialization progress on entomopathogenic nematodes in China

YAN Xun^{1**} GUO Wen-Xiu² TIAN Cheng-Li^{3,4} LI Xing-Yue⁵ CHEN Shu-Long⁶
JIN Yong-Ling⁷ LI Li-Ying⁸ TANG Rui⁸ CAO Li^{8***} HAN Ri-Chou^{8***}

(1. Key Laboratory of Green Prevention and Control on Fruits and Vegetables in South China, Ministry of Agriculture and Rural Affairs, College of Agriculture & Biology, Zhongkai University of Agriculture and Engineering, Guangzhou 510225, China; 2. Shandong Key Laboratory for Green Prevention and Control of Agricultural Pests, Institute of Plant Protection, Shandong Academy of Agricultural Sciences, Jinan 250100, China; 3. Institute of Agricultural Resource and Environment, Jilin Academy of Agricultural Sciences (Northeast Agricultural Research Center of China), Changchun 130033, China; 4. College of Biological Sciences, Liaoning University, Shenyang 110136, China; 5. Key Laboratory of Integrated Pest Management on Crops in Southwest, Ministry of Agriculture and Rural Affairs, Institute of Plant Protection, Sichuan Academy of Agricultural Sciences, Chengdu 610066, China; 6. Institute of Plant Protection, Hebei

*资助项目 Supported projects: 国家重点研发计划“政府间国际科技创新合作”重点专项(2024YFE0109600); 广东省科学院发展专项资金项目(2022GDASZH202201010); 国家自然科学基金面上项目(32172491); 国家自然科学基金青年项目(32302452)

**第一作者 First author, E-mail: yanxun@zhku.edu.cn

***共同通讯作者 Co-corresponding authors, E-mail: caol@giz.gd.cn; hanrc@giz.gd.cn

收稿日期 Received: 2025-08-12; 接受日期 Accepted: 2025-09-10

Academy of Agriculture and Forestry Sciences, Baoding 071000, China; 7. Heilongjiang Bayi Agricultural University, Daqing 163319, China; 8. Guangdong Key Laboratory of Animal Conservation and Resource Utilization, Guangdong Public Laboratory of Wild Animal Conservation and Utilization, Institute of Zoology, Guangdong Academy of Sciences, Guangzhou 510260, China)

Abstract Entomopathogenic nematodes (EPNs), as parasitic natural enemies of insects that have been commercially applied internationally and are highly effective against soil-borne and both-boring pests, have played a significant role in the green management of important pests in agriculture, forestry, and health since their introduction to China since the 1960s. Globally, over 150 species of EPNs have been isolated and identified. In China, 27 species of *Steinernema* and 6 species of *Heterorhabditis* have been discovered, including 19 new species of *Steinernema* and 2 new species of *Heterorhabditis*. Research on the symbiotic bacteria of EPNs has clarified the symbiotic specificity between these bacteria and the nematodes. Numerous toxins and secondary metabolites with insecticidal, antibacterial, and nematocidal activities have been identified from these symbiotic bacteria, demonstrating broad application prospects. The effectiveness of EPNs in field applications is influenced by environmental factors. Studies on the environmental tolerance and underlying mechanisms of EPNs have led to the selection of strains adapted to environmental pressures, which are being used to control specific pests in different regions. Research and technological innovations in the cultivation, formulation, preservation, and application techniques of EPNs have enabled their industrial production in China. Currently, EPNs have been widely adopted in China for pest control, achieving remarkable results in managing pests such as the peach fruit moth (*Carposina nipponensis*), grubs, and chive maggots (*Bradysia odoriphaga*) in Chinese chives. This paper comprehensively reviews the research and development progress of these nematodes in China on their identification and classification, industrialization and formulation, symbiotic bacteria, association with the symbiotic bacteria, applications, environmental tolerance, and the relationship with abiotic environmental factors.

Key words entomopathogenic nematodes; symbiotic bacteria; identification and classification; commercial production; formulation; field application

斯氏线虫科 *Steinernematidae* 和异小杆线虫科 *Heterorhabditidae* 昆虫病原线虫是目前国际上对土栖性和钻蛀性害虫特效的天敌类群。这类线虫以 3 龄感染期幼虫 (Infective juveniles, IJ) 主动寻找昆虫寄主, 对非靶标生物和生态环境安全, 可离体产业化生产, 已广泛用于防治多种害虫 (Ehlers, 2001; Gaugler *et al.*, 2002; 吴文丹等, 2014; Labaude and Griffin, 2018; 王杰等, 2021), 德国 E-nema、荷兰 Koppert、英国 Becker Underwood、日本 SDS 以及我国潍坊宏润农业科技有限公司、河南济源生物科技有限公司等多家公司企业均商业化生产这类生物农药 (Han *et al.*, 2000; Kaya *et al.*, 2006; Peters, 2013; Shapiro-Ilan *et al.*, 2017; 王杰等, 2021)。

利用线虫防治有害昆虫的研究是 20 世纪 30 年代才开始的。我国从 20 世纪 60 年代开始昆虫病原线虫研究和应用, 最早由中国科学院动物研究所沙搓云先生从捷克引入昆虫病原线虫。自

1978 年广东省科学院动物研究所 (时称广东省昆虫研究所) 所长李丽英研究员先后从澳大利亚、美国和加拿大引进了多种昆虫病原线虫, 率先在我国开展昆虫病原线虫的大量繁殖与田间应用研究。从 1986 年起, 广东省科学院动物研究所和中国农业科学院植物保护研究所 (当时为中国农业科学院生物防治研究所) 与澳大利亚 CSIRO 组织的昆虫学研究所共同实施长达 12 年的中澳国际合作项目, 广东省科学院动物研究所与德国基尔大学、爱尔兰国立大学以及印度尼西亚 The University of Jember 共同承担欧盟第五框架国际合作项目。在国内外项目的资助下, 我国昆虫病原线虫研究和产业化走进了新时代, 昆虫病原线虫在农林和卫生等害虫的可持续治理中发挥了巨大的作用。

本文从昆虫病原线虫的鉴定和分类、产业化、共生细菌、环境耐受性和应用, 昆虫病原线虫与非生物环境因子等领域全面综述我国对这一天敌种类的研发进展。

1 昆虫病原线虫的鉴定和分类

目前已从全球分离和鉴定了 120 种斯氏属 *Steinernema* 线虫和 30 种异小杆属 *Heterorhabditis* 线虫 (Kozodoi, 1984; Gardner *et al.*, 1994; 刘杰, 1994; Stock, 1996; Stock *et al.*, 1996, 2009; Ganguly and Singh, 2000; Cutler and Stock, 2003; Plichta *et al.*, 2009; Clausi *et al.*, 2011; Gorgadze *et al.*, 2015; Shahina *et al.*, 2017; Bhat *et al.*, 2020, 2023a, 2023b; Didiza *et al.*, 2021; Lis *et al.*, 2021; Machado *et al.*, 2022, 2025; Tian *et al.*, 2022; Patil *et al.*, 2023; Soni *et al.*, 2023; Baniya *et al.*, 2024; Anes *et al.*, 2025; Zhan *et al.*, 2025; NCBI 数据库 <https://www.ncbi.nlm.nih.gov/taxonomy/?term=Steinernema> 和 <https://www.ncbi.nlm.nih.gov/taxonomy/?term=heterorhabditis>)。

我国气候、植被和生态系统的多样性使得昆虫病原线虫资源丰富。李小峰和王国汉 (1989) 调查结果表明, 福建、广东、海南等地均有斯氏线虫和异小杆线虫的分布; 陈书龙等 (2006) 调查河北省线虫资源, 分离出斯氏线虫 17 个种群, 异小杆线虫 49 个种群; Ma 等 (2010) 调查了中国北方线虫资源, 共获得斯氏线虫 7 个种, 103 个品系, 异小杆线虫 3 个种, 102 个品系; 钱秀娟等 (2014) 获得 29 个线虫品系, 其中 *S. affine*、*S. karil* 和 *S. kraussei* 为国内新记录种; Wang 等 (2014) 对辽宁省昆虫病原线虫资源进行调查, 发现新记录种 *S. bicornutum*; 张红玉 (2016) 调查山东地区线虫资源, 获得 6 个品系; 王思雨 (2022) 对华北北部地区进行调查, 获得 53 个品系; 刘佳等 (2025) 补充调查了甘肃昆虫病原线虫的资源情况, 共获得 20 个品系。

目前, 我国发现斯氏线虫共 27 种, 其中新种 19 种, 分别是 *Steinernema caudatum* (Xu *et al.*, 1991)、*Steinernema longicaudum* (Shen and Wang, 1992)、*Steinernema ceratophorum* (Jian *et al.*, 1997)、*Steinernema guangdongense* (Qiu *et al.*, 2004)、*Steinernema aciari* (Qiu *et al.*, 2005c)、*Steinernema akhursti* (Qiu *et al.*, 2005a)、*Steinernema beddingi* (Qiu *et al.*, 2005b)、

Steinernema sichuanense (Mráček *et al.*, 2006)、*Steinernema hebeiense* (Chen *et al.*, 2006a)、*Steinernema leizhouense* (Nguyen *et al.*, 2006)、*Steinernema cholashanense* (Nguyen *et al.*, 2008a)、*Steinernema xueshanense* (Mráček *et al.*, 2009)、*Steinernema pui* (Qiu *et al.*, 2011)、*Steinernema changbaiense* (Ma *et al.*, 2012a)、*Steinernema xinbinense* (Ma *et al.*, 2012b)、*Steinernema tielingense* (Ma *et al.*, 2012c)、*Steinernema taiwanensis* (Tseng *et al.*, 2018)、*Steinernema populi* (Tian *et al.*, 2022) 和 *Steinernema tarimense* (Zhan *et al.*, 2025)。

我国报道的异小杆线虫种类较少, 共 6 种, 包含 2 个新种 (其中 1 种为同种异名)。Hsieh 等 (2009) 在中国台湾地区发现 *Heterorhabditis brevicaudis*, 并补充分子描述, Hunt and Subbotin (2016) 认为 *H. brevicaudis* 和 *H. indica* 是同种异名。Li 等 (2012) 在北京樱桃树下发现昆虫病原线虫新种 *Heterorhabditis beicherriana*。异小杆线虫 *Heterorhabditis marelatus*、*H. bacteriophora*、*H. indica* 和 *H. megidis* 在我国均有分布, 其中 *H. megidis* 分布范围最广, 是主要的优势种群。Nguyen 等 (2008b) 基于系统进化分析, 将异小杆属划分为 3 个分支: “*Indica*”、“*Bacteriophora*”和 “*Megdis*”。我国发现的异小杆线虫属于 “*Indica*” 分支的为 *H. indica*; 2 种属于 *Bacteriophora* 分支: *H. beicherriana* 和 *H. bacteriophora*; 2 种属于 *Megdis* 分支: *H. marelatus* 和 *H. megidis*。

2 昆虫病原线虫的产业化

我国科研人员在昆虫病原线虫的培养、剂型配制及保存等方面开展了大量的研发, 提高了昆虫病原线虫生产效率, 延长货架时间, 提高田间防效, 降低使用成本。

2.1 昆虫病原线虫产业化培养

昆虫病原线虫的产业化培养方法包括昆虫活体培养及离体培养, 离体培养则包括固体培养

和液体培养两种方法。在不同培养方法的工艺优化上开展的技术改进,为实现昆虫病原线虫的产业化培养和应用提供了有力的支撑(颜珣和韩日畴, 2016)。

常用于昆虫病原线虫活体培养的昆虫包括大蜡螟 *Galleria mellonella*、黄粉虫 *Tenebrio molitor*、家蚕 *Bombyx mori*、适宜特定昆虫病原线虫品种的金龟子或其他直翅目昆虫等(Peters *et al.*, 2017; Han *et al.*, 2024; 田成丽, 2024)。徐洁莲和谢汝创(1992)比较了 *S. carpocapsae* 和 *S. anomali* 两种线虫在大蜡螟和鸡杂海绵培养基下连续传代培养后的差异。李春杰等(2008)使用繁殖袋,筛选蛭石作为培养介质,以末龄黄粉虫幼虫为活体寄主进行 *H. bacteriophora* Hb 的大量繁殖。阎爱华等(2015)研发了实验室内活体大量繁殖昆虫病原线虫的装置。姚红燕等(2018)以较易获得的甜菜夜蛾 *Laphygma exigua* 高龄幼虫为寄主,有效降低昆虫病原线虫活体繁殖的成本。李小龙等(2019)研发了智能一体化昆虫病原线虫活体培养装置,减少人力,提高了培养效率。*H. megidis* 0627M 和 *S. feltiae* 0619HT 两种品系可以通过寄主韭蛆复壮,使其生殖力、致病力和侵入率都明显升高(狄斌等, 2021)。

共生细菌对昆虫病原线虫单菌培养发挥重要作用。足量的共生细菌才能提高昆虫病原线虫的产量(徐洁莲, 1989; 杨秀芬和杨怀文, 1997)。固体培养系统中,初生型菌比次生型菌培养线虫的产量提高 3-10 倍(韩日畴等, 1990a)。次生型共生细菌未被 *H. bacteriophora* 线虫携带,也不能支撑线虫生长(Han and Ehlers, 2001)。*P. luminescens* 初生型菌产生的细胞间晶体蛋白 CipA 和 CipB 可以诱导 *Steinernema* 线虫恢复、发育和繁殖(You *et al.*, 2006)。韩日畴等(1990b)报道了通过消毒昆虫病原线虫卵获得无菌幼虫的方法。无菌 *H. bacteriophora* IJ 可在被感染的无菌大蜡螟幼虫血淋巴中恢复发育到 1 龄幼虫,但未能杀死大蜡螟幼虫,而无菌 *S. carpocapsae* 能够致死无菌大蜡螟幼虫并产生后代,但 IJ 产量低于有共生细菌存在时的产量(Han and

Ehlers, 2000)。丘雪红等(2004)发现共生细菌除了显著影响 IJ 的致病力,也影响线虫干重、蛋白质、氨基酸、糖原和脂肪酸等生化物质含量。

在建立昆虫病原线虫体外单菌培养技术的基础上,通过优化培养基组分、培养设备、培养流程等,提高体外培养的效率。最早使用的线虫固体培养基是从澳大利亚引进的鸡杂海绵培养基(Bedding, 1981),早期的固体培养基主要包含猪杂/鸡杂、水、猪油/牛油等简单的成分,混合海绵配制而成(赵忠懿等, 1991)。李素春等(1987)最早研究了培养基成分、培养基含水量、线虫接种量、共生细菌接入量及培养温度等因素对引进的 *Neoapectana feltiae* 线虫在鸡杂海绵培养基中产量的影响。杨怀文等(1991)以容易获得的鸭肠、油脂和柞蚕 *Antheraea pernyi* 蛹配成培养基培养 *S. feltiae* 线虫,产量与鸡内脏配方相当。Han 等(1992)报道,添加玉米粉、鱼粉等含脂量高的成分可以提升 *Steinernema* sp. CB2B 和 *H. bacteriophora* H06 固体培养产量。Han 等(1993)以线虫接种量、温度和培养时间为参数,利用正交旋转组合设计方法优化了 *S. carpocapsae* 线虫的固体培养系统。潘洪玉等(1995)筛选利用植物蛋白质替代鸡肠的配方,以低成本获得稳定的线虫产量。韩日畴等(1995)优化了固体干粉培养基组分,获得 *S. carpocapsae* A24 和 *H. bacteriophora* H06 线虫最高产量达 5.8×10^5 和 4.8×10^5 IJ/g,并在此基础上制定线虫工厂化固体培养标准化流程和质量控制程序(韩日畴等, 1997)。于固体培养基中加入大蜡螟幼虫组织匀浆,可使出现衰退的 *S. carpocapsae* Beijing 线虫恢复繁殖力(常韶华和杨怀文, 1998)。王忠文等(1999)用鸡蛋替代鸡杂,固体培养可收获 6.07×10^5 IJ/g,发育历期最短,3-4 d 便可增殖一代。郝杰等(2017)对 *S. carpocapsae* HB310 线虫的固体培养基及培养条件进行优化,筛选到更为经济、高效的线虫固体培养基配方。Zhen 等(2018)比较了固体培养基中添加大蜡螟、黄粉虫和丝光绿蝇 *Lucillia sericata* 昆虫干粉对 *S. feltiae* SN 线虫产量的影响,发现添加 5% 黄粉虫干粉的产量最高,生产的 IJ 对斜纹夜蛾

Spodoptera litura 的致死速度也最高。Wang 等 (2022) 研究发现, 线虫信息素蛔苷 (Ascarosides) 可显著提高昆虫病原线虫的产量。赵国玉和郭文秀 (2023) 在 *S. glaseri* 的固体培养基中添加黄粉虫干粉、蝇蛆干粉等成分并优化培养条件, 可使线虫最高产量提高 47.5%, 培养周期缩短 46.4%, 解决了这一线虫培养周期长、产量低、培养成本高的问题。

液体大量培养技术是昆虫病原线虫的商品化发展方向。首先在摇床中建立了 *S. carpocapsae* A24 和 *H. bacteriophora* H06 线虫的液体培养, 线虫接种量和培养时间显著影响线虫产量 (Han, 1996)。朱明军等 (1998) 在摇瓶培养的基础上, 在国内率先开始了应用发酵罐进行昆虫病原斯氏线虫 A24 品系的液体培养, 发现气升内环流反应器最适合昆虫病原线虫的培养, 优化了液体培养条件, 为反应器大规模培养线虫提供了依据 (朱明军等, 1999)。站丽莉等 (2008) 优化了 *S. carpocapsae* 线虫 2 种液体培养基的产量和效率。王丽芳等 (2010) 比较了 *H. bacteriophora* Hb-1 的液体和固体繁殖技术, 优选了线虫培养基。李春杰等 (2007) 比较了固体、液体等 4 种不同方法培养 *H. bacteriophora* Hb-1 和 *S. carpocapsae* Sc-2 的效果。

在专利申报中, 韩日畴和李丽英 (2003) 在昆虫病原线虫液体单菌培养系统中, 于线虫发育的关键龄期加入 0.01-1 $\mu\text{g/mL}$ 的链霉素, 以改变菌体密度和诱导共生菌产生变异型, 进而控制 IJ 的形成, 提高 IJ 的产量。杨怀文等 (2007) 利用淀粉酶、糖化酶、蛋白酶和脂肪酶等酶制剂对线虫培养基进行酶解处理后培养昆虫病原线虫, 线虫产量比未酶解培养基提高 40%, 收获期提前 4-6 d。简恒等 (2011) 将昆虫酶解物作为组份加入到培养基中, 可以提高线虫产量、缩短培养周期、降低生产成本。刘勇 (2020) 选用 20% (m/v) 面包虫和 4% (m/v) 玉米粉作为基本培养基, 线虫繁殖量达 2.6×10^7 IJ/瓶; 添加 2% (m/v) 活性炭和 3% (m/v) Fe_3O_4 能有效提高繁殖率。Kong 等 (2022) 和谷星慧等 (2023) 发现在培养基中添加二甲基亚砷 (Dimethyl sulfoxide, DMSO)

及蛔苷可显著提高 *S. carpocapsae* All 和 *H. bacteriophora* H06 的 IJ 产量。DMSO 主要通过抑制 *Hint module*、*PAN domain protein*、*Dpy-13* 3 个基因来调控线虫的怀卵量, 从而提高 IJ 产量 (Liu et al., 2024)。

广东省科学院动物研究所 (原广东省昆虫研究所) 改进的昆虫病原线虫固体培养技术已在多家公司转化, 实现 *S. carpocapsae*、*S. feltiae*、*S. longicaudum*、*H. bacteriophora*、*H. indica* 等线虫的规模化生产及商品化, 并通过技术转移及培训, 在印度尼西亚、巴基斯坦、卢旺达、朝鲜、韩国等国家成功应用, 改变了这些国家的害虫防治模式, 在降低化学农药依赖、促进环境可持续发展方面成效显著 (王杰等, 2021; Yan et al., 2025)。此外, *S. feltiae* A24, *S. bibionis* T319 和 *S. glaseri* NC34 (杨平和徐洁莲, 1989)、*S. feltiae* OTIO (王慧等, 1998)、异小杆线虫 NLK-1 (布卡·欧尔娜等, 2015)、*S. ceratohporum* (Yan et al., 2021)、*S. glaseri* (赵国玉和郭文秀, 2023) 等昆虫病原线虫品种和品系也已实现大量培养, 用于田间害虫防控。

培养设备改进对昆虫病原线虫体外培养流程的优化同样具有重要意义。孙瑞红和李爱华 (2011) 设计了带有清洗筛的线虫清洗器。田会鹏等 (2013) 设计了 *S. carpocapsae* A24 的不锈钢培养盒。许艳丽等 (2018) 发明了一种可以同时进行灭菌和搅拌的装置, 搅拌桨以电机驱动, 相比人工操作简单, 该装置用于昆虫病原线虫固体繁殖。郭文秀等 (2019d, 2019e) 改进了昆虫病原线虫的培养盒, 以简单的结构和低加工成本实现透气保湿环境及昆虫病原线虫的自动化洗出分离。陈瑞等 (2020) 采用耐高温塑料材质设计了新型培养瓶。李小龙等 (2021) 设计使用培养袋培养昆虫病原线虫。廖慧君等 (2023a) 发明了通过固定组件和隔断组件的配合, 将长条状海绵经隔断组件和固定组件后被粉碎成海绵块的装置, 用于昆虫病原线虫培养基混合粉碎, 操作简单, 自动化高, 实用性强。廖慧军等 (2023b) 还发明了一种发酵菌种液体培养系统, 解决了外部空气与发酵罐内部温差过大的问题。阮维斌

(2023)研发了利用挤压板运动挤压海绵分离线虫的装置。

2.2 昆虫病原线虫剂型、保存及施用技术

徐洁莲和刘秀玲(1997)研究昆虫病原线虫常温保存技术。韩日畴等(2000)筛选出国产硅藻土可以用于 *S. carpocapsae* 线虫的贮存,添加食品抗霉剂能维持 2 周的防霉效果。Feng 等(2006)筛选了甘油和加强型人工海水配制的渗透压溶液,能够诱导 *S. carpocapsae* All 线虫进入脱水休眠状态,进而提高线虫耐热性,延长保存时间。Yan 等(2010)在此基础上研究了该渗透压溶液对 15 个不同品种和品系线虫的作用效果,渗透压处理后线虫耐热性显著提高,为应用渗透压溶液诱导昆虫病原线虫进入脱水休眠状态进而长期保存线虫提供了基础。李春杰等(2014)研究了 *H. bacteriophora* HBN 在 4 种介质(蛭石、海绵、沙土和珍珠岩)中冷冻储存的最优条件;进一步研究发现逐渐降温预处理可以显著提高 *H. bacteriophora* HBN 和 *H. bacteriophora* NJ 两种线虫的冷冻存活率(李春杰等,2015)。马小雅(2016)发明了贮存线虫的多种保存剂。肖宇辉(2018)报道了海绵保存 *S. costaricense* 线虫的保水剂、防腐剂和洗出剂。郭文秀等(2020)发明了利用腐殖土保存多种昆虫病原线虫的方法。

剂型研发方面,也通过改进制剂配方,配制剂型延长昆虫病原线虫货架时间,同时针对不同的目标害虫研发适合田间应用的剂型。徐洁莲和杨平(2000)研究发现国产水溶性粘合剂 POA 对 *S. carpocapsae* A24 有增效的效果。陈松笔等(2001c)筛选了含有 10% CYL、1% 羧甲基纤维素钠和 0.01% 4203 的蒸发抑制剂、保湿剂和乳化剂组合,作为 *S. carpocapsae* CB-16 线虫叶面用抗干燥剂。张安邦(2015)提供了 *S. glaseri* Sgib 的粉剂配方(含 30% 白炭黑、69.4% 水、0.5% 羧甲基纤维素钠和 0.1% 甲醛),该粉剂在 10 °C 储存 30 d 后线虫的存活率为 100%,储存 60 d 线虫存活率为 92.38%。Guo 等(2017b)筛选出 1:0.8-1:1.2 (w/w) 的蛭石和腐殖土适合作为载体

保存线虫,添加 0.1%-0.2% 的杀菌剂 Proxel GXL 可以延长线虫保存时间,保存过程中 IJ 的感染力不受影响。方分分等(2019)以蔗糖、葡萄糖、海藻糖、山梨醇等作为稳定剂(0.2%-1.0% 质量比),以甘油、黄原胶、羧甲基纤维素等为保水剂(0.2%-1.0% 质量比),以山梨酸钾、丙酸钠、富马酸二钾酯、甲醛等为防霉剂(0.1%-0.5% 质量比),以硅藻土、白炭黑、可溶性淀粉、天然沸石粉等作为载体(40%-50% 质量比),配制的昆虫病原线虫粉剂,在 10 °C 条件下贮存 3 个月后线虫存活率 80% 以上。李星月等(2020)将 *H. beicheriana* 与硫脲复配成的制剂,含有有机硅、矿源黄腐酸钾、海藻糖及共生细菌培养基等成分,防治花生蛱蛭及其他地下害虫防效达 60% 以上。南宫自艳等(2022)以白炭黑和/或粘土为基质,以羟丙基甲基纤维素和/或羧甲基纤维素钠为保水剂,以海藻酸钠和/或黄原胶为紫外保护剂,以山梨酸钾和/或甲醛为防腐剂,加入甘油和水配制的线虫粉剂,能显著提高线虫长时间贮存后的存活率。李晶晶等(2023)混合黄粉虫、木屑、水并接种线虫,培养获得黄粉虫虫尸,可直接作为昆虫病原线虫制剂应用田间。刘琴等(2023)以草地贪夜蛾 *Spodoptera frugiperda* 的蛹生产 *S. carpocapsae*,形成蛹蜕自然包裹线虫的蛹胶囊,再以 10% 聚乙烯醇溶液(w/w)喷淋蛹体,在蛹表面形成水溶性半透膜,获得斯氏线虫蛹胶囊制剂,在 20-25 °C 保存,线虫存活率仍高达 78% 以上。郭文秀等(2024)研发了一种保湿缓释胶囊,可以保护昆虫病原线虫不受土壤干燥的影响。李星月等(2025)研发了包括昆虫病原线虫 *S. carpocapsae* 黄粉虫虫尸剂、甘薯小象甲 *Cylas formicarius* 人工饲料、多酚氧化酶抑制剂和海藻糖干粉的虫尸复配剂。

采用喷雾、滴注、挂物和涂布等不同方法施用格氏线虫 85011 品系防控直锥大象虫 *Cyrtorachelus longimanus* 时,发现在淀粉糊中加线虫做成糊剂涂布竹笋虫蛀口的方法防效最高(刘南欣等,1990)。以高压机动喷雾机施用昆虫病原线虫防控桃小食心虫 *Carposina nipponensis*,压力在 10-20 kg/cm² 时,对线虫的

存活、寄主的侵染力和寄生强度均无明显影响(黎彦等, 1991)。陈镜华和韩日畴(2014)发明了昆虫病原线虫喷施设备。郭文秀等(2019f)、廖慧军等(2023c)发明的储存装置, 可以使水体中的昆虫病原线虫保持悬浮状态, 解决昆虫病原线虫施用过程中水中存活时间较短的问题。郭文秀等(2019a, 2019b)发明了昆虫病原线虫田间随水施加装置和可切换施用不同昆虫病原线虫的田间施用装置。郭文秀等(2019c)还研发了用于防治天牛的昆虫病原线虫注射装置, 能够在田间防治天牛时实现连续定量注射。这些设备的研发, 提升了田间作业效率, 使昆虫病原线虫施放均匀, 节省了时间, 降低了人工成本和劳动强度, 还相应提升了昆虫病原线虫的防治效果。

3 昆虫病原线虫-细菌共生体

昆虫病原线虫 IJ 体内携带有互利共生的共生细菌。昆虫病原线虫通过寄主的口、肛门、气孔等自然开口、伤口或体壁节间膜进入昆虫体腔内, 释放共生细菌到寄主血淋巴中, 共生细菌在昆虫病原线虫生活史中发挥重要作用: 共生细菌在寄主体内繁殖, 分泌多种毒素, 使害虫患败血症死亡; 同时产生多种胞外酶, 将昆虫物质转化为能被线虫所吸收利用的营养物质, 支持线虫生长; 共生细菌还分泌抗菌物质抑制非共生细菌的生长, 为线虫生长发育提供稳定环境(吴文丹等, 2014; 王杰等, 2021)。昆虫体内营养耗尽后, 昆虫病原线虫-细菌共生体重新形成, 离开虫体, 寻找新的寄主。

3.1 共生细菌的种类

隶属于变形菌门 Proteobacteria 的 γ -亚纲肠杆菌科 Enterobacteriaceae 的致病杆菌属 *Xenorhabdus* 和发光杆菌属 *Photorhabdus* 细菌, 革兰氏阴性, 不产孢, 兼性厌氧, 具有支持线虫的共生性与致死昆虫的病原性特点(Eckstein et al., 2019; Stock, 2019)。*Xenorhabdus* 和 *Photorhabdus* 属共生细菌分别与斯氏线虫属和异小杆线虫属线虫共生。

目前已确定的 *Xenorhabdus* 属细菌有 27 种,

Photorhabdus 属细菌有 19 种(Machado et al., 2021)。2003 年, 第一版 *P. luminescens* 基因组公布(Duchaud et al., 2003), 随后有多个共生细菌的基因组公布(Ghazal et al., 2017; Duong et al., 2019; Somvanshi et al., 2019), 包括我国学者测序的多个共生细菌基因组(Qiu et al., 2014; Li et al., 2021a)。

3.2 共生细菌的特征

Xenorhabdus 和 *Photorhabdus* 属共生细菌在体外培养时出现“初生型”和“次生型”的菌落和菌体(Akhurst, 1980)。能被 IJ 携带的仅为初生型细菌。初生型细菌更好地支持昆虫病原线虫生长和发育, 次生型细菌会影响昆虫病原线虫产量(Akhurst and Boemare, 1990; Han and Ehlers, 2001)。初生型 *P. luminescens* 细菌产生两种胞内晶体蛋白 CipA 和 CipB, 而次生型缺少晶体蛋白; 胞内晶体蛋白被证明为线虫繁殖的营养因子(You et al., 2006)。

3.3 共生细菌的作用

我国学者发现, 共生细菌培养液对昆虫、植物病原或植物线虫具有控制作用。*P. luminescens*、*X. nematophila* A24-1 和 *X. nematophila* HB310 菌液分别对甜菜夜蛾(王欢等, 2007)、东亚飞蝗 *Locusta migratoria manilensis*(张园等, 2013)、小菜蛾 *Plutella xylostella*(Wu et al., 2020)具有致死作用。共生细菌菌液对大蜡螟幼虫(付俊瑞等, 2018a, 2018b)、小菜蛾幼虫(张红玉等, 2016)和南方根结线虫 *Meloidogyne incognita* 卵的孵化和幼虫具有抑制作用(王鑫鹏, 2018; 吴黎黎等, 2024)。*Xenorhabdus* 和 *Photorhabdus* 菌株分别抑制番茄灰霉菌 *Botrytis cinerea*(刘霞等, 2006)、苹果灰霉菌 *B. cinerea*(文旭等, 2012)、葡萄灰霉病(曹林青等, 2021)、辣椒疫霉菌 *Phytophthora capsici*(王永宏和张兴, 2007)。共生细菌可控制玉米小斑病菌 *Bipolaris maydis*(王欢等, 2009)、串珠镰刀菌 *Fusarium moniliforme*(崔安妮等, 2015)、黑曲霉 *Aspergillus niger*(高宇洁等,

2022)、草莓炭疽菌 *Colletotrichum fragariae*、稻瘟病菌 *Magnaporthe oryza*、大豆疫霉菌 *Phytophthora sojae*、尖孢镰刀菌 *Fusarium oxysporum* 和辣椒立枯丝核菌 *Rhizoctonia solani* (张金花等, 2024) 等病菌引起的植物病害。

共生细菌产生的多种代谢产物应用潜力巨大 (Tobias *et al.*, 2017; Dreyer *et al.*, 2018; Shi *et al.*, 2019; Neubacher *et al.*, 2020; Meesil *et al.*, 2025)。不同种类或菌株产生功能和作用效果不同的多种代谢物活性成分 (方香玲等, 2008; Liu *et al.*, 2019; Mollah *et al.*, 2020; Wang *et al.*, 2022)。*Xenorhabdus* 和 *Photorhabdus* 共生细菌分泌的各种抗菌物质已用于研发新型抗菌素 (Shi and Bode, 2018; Muangpat *et al.*, 2020)。我国学者在这一领域做出了重要贡献, 比如从共生细菌中分离了多种抗菌活性物质, 包括 Xenocoumacin 1 和 Xenocoumacin 2 (Guo *et al.*, 2017a; Zhang *et al.*, 2024)、Xenematide G (Xi *et al.*, 2019)、PAX1 肽与异香豆素 (Dreyer *et al.*, 2019)、Nematophin (Zhang *et al.*, 2019a)、Darobactin (Imai *et al.*, 2019)、环肽 Photoditritide (Zhao *et al.*, 2019)、非核糖体肽 (Du *et al.*, 2022) 以及多种挥发性化合物 (Baïome *et al.*, 2022)。pH 调节 (Guo *et al.*, 2017a)、原位产品移除法 (Dong *et al.*, 2020) 和发酵过程优化可提高 Xenocoumacin 1 的产率 (Han *et al.*, 2024)。由共生细菌从吩嗪 Phenazine 转化的抗菌素, 有利于维持线虫-细菌复合体的特殊环境生态位 (Shi *et al.*, 2019)。此外, *X. bovienii* 通过合成放线菌素 D (Actinomycin D) 诱导大蜡螟幼虫细胞凋亡和败血症 (Zhang *et al.*, 2023)。蒽醌可通过大肠杆菌 *Escherichia coli* 表达 *P. luminescens* 的 *antABCDEFGH* 基因簇获得 (Zhou *et al.*, 2019)。大肠杆菌通过整合 *Photorhabdus* 属细菌 type II polyketide synthase 基因可大量生产胭脂红酸 (Yang *et al.*, 2021)。目前被认为极具商业价值的抗癌药物 Glidobactin A 和 Cepafungin I 便来源于共生细菌 (Zhao *et al.*, 2020)。于 *P. laumondii* 发现可用于抑制肿瘤的

Glidobactin 样蛋白酶体抑制剂 (Zhao *et al.*, 2021)。*X. stockiae* HN_xs01 菌株被证明具有抗肿瘤作用 (Zhang *et al.*, 2022)。共生细菌分泌抗菌物质的调控机制研究方面, *cpxR* 基因被认为影响 *X. nematophila* 抗菌物质合成 (Guo *et al.*, 2019; Zhang *et al.*, 2019b)。*OmpR* 被证明间接调控 *X. nematophila* 菌株中 Xenocoumacin 1 的合成 (Han *et al.*, 2025)。产生 Odilorhabdins 抗菌素的共生细菌具有自我保护机制, 编码 N-乙酰转移酶的基因 *oatA* 负责 Odilorhabdins 抗性 (Lanois-Nouri *et al.*, 2022)。

共生细菌产生的各类毒性蛋白可有效作用于不同种类昆虫、植物线虫和原生动物 (Bowen *et al.*, 2000; Pan *et al.*, 2002; Tobias *et al.*, 2018; Muangpat *et al.*, 2020)。除了 *Photorhabdus* insect related protein (Pir) (Waterfield *et al.*, 2005), Makes Caterpillars Floppy (Mcf) (Daborn *et al.*, 2002), Photox (Visschedyk *et al.*, 2012), 几丁质酶 (Liu *et al.*, 2019; Mahmood *et al.*, 2020) 和 Galtox (Ahuja *et al.*, 2021), 我国学者还发现了可作为强效的蛋白传递系统 (Jiang *et al.*, 2022) 的 *Photorhabdus* virulence cassettes (PVC) (Yang *et al.*, 2006; Wang *et al.*, 2022, 2024)、*Photorhabdus* insecticidal toxin (Pit) (Li *et al.*, 2009) 和 PaxAB (Zhang *et al.*, 2014)。证明 *X. nematophila* HB310 菌株的几丁质酶 *chi60* 和 *chi70* 基因对 Tc 毒性是必须的 (Liu *et al.*, 2022), N-聚糖和硫酸化糖胺聚糖介导 Tc 对哺乳动物细胞的毒性 (Song *et al.*, 2021)。这类细菌产生的胞内和胞外蛋白可抑制棉铃虫 *Helicoverpa armigera* 初孵幼虫 (杨保军等, 2007)。共生细菌中分离获得的非核糖体肽 Rhabdopeptides, 可用于拮抗携带热带疾病病原体的原生动物 (Wesche *et al.*, 2018), 致死南方根结线虫 *M. incognita* (Bi *et al.*, 2018)。我国学者从 *X. innexi* DSM 16336 中获得了 7 个 Rhabdopeptide/xenortide 肽, 经测定, 这些多肽对布氏罗得西亚锥虫 *Trypanosoma brucei* 和恶性疟原虫 *Plasmodium falciparum* 产生抑制作用 (Zhao

et al., 2018), 为新药物研发开辟途径。发光杆菌产生的 *TcaB*、*Txp40*、*PaTox*、*PirAB*、*PTC* 等毒素对昆虫具有口服或注射活性 (Wu and Yi., 2016; Bogdanovic et al., 2019; Mathur et al., 2019; Nangong et al., 2020; Shankhu et al., 2020; Wang et al., 2023)。致病杆菌产生的 *Xax*、*srfABC*、*PirAB*、*Tp40*、*hipBA* 等毒素也具有强烈的杀虫活性 (杨君等, 2008; 郭笑笑等, 2019; Yang et al., 2019; Yadav and Rathore, 2020)。X. stockiae 菌株产生的 *SrfABC* 毒素对棉铃虫幼虫呈现杀虫活性, 推测在宿主定殖中起重要作用 (Yang et al., 2019), 并表现细胞毒性和导致细胞凋亡 (Sun et al., 2019)。目前已从大蜡螟幼虫中获得与 X. nematophila *PirAB* 有交互作用的 Arylphorin 蛋白 (Nangong et al., 2020)。转 *TcdA* 基因的拟南芥 Arabidopsis thalian 对烟草天蛾 Manduca sexta 和黄瓜点叶甲 Diabrotica undecimpunctata 幼虫具有毒杀作用 (Liu et al., 2003)。转 P. luminescens 杀虫基因 *TcdA1* 和 *TcdB1* 的阴沟肠杆菌 Enterobacter cloacae 对台湾乳白蚁 Coptotermes formosanus (Zhao et al., 2008) 和斜纹夜蛾、甜菜夜蛾、黄曲条跳甲 Phyllotreta striolata 均具有杀虫活性 (张鹏波等, 2010)。P. temperata Meg1 基因簇 *pttABC* 在大肠杆菌中表达可获得一种新的环脂肽 Phototemtide A, 对恶性疟原虫具有选择性抗原动物活性 (Zhao et al., 2019, 2020)。

前期研究发现, 热灭活的 P. luminescens TT01 菌体 (Wu et al., 2014) 或脂多糖 (Wu et al., 2015a) 诱导大蜡螟幼虫免疫致敏, 但 *PirA2B2* 蛋白免疫致敏的大蜡螟幼虫对 P. luminescens TT01 的感染没有明显的抵抗 (Wu et al., 2015a)。胰岛素促进共生细菌的生长发育 (杨亚贤等, 2023; 李茜童等, 2024; 钱秀娟等, 2024)。T3SS 和鞭毛都促进了共生细菌在宿主细胞内的侵袭和定植, 对次生代谢产物的形成和分泌影响较小 (Huang et al., 2024)。

基于来自 P. luminescens 的寄主特异性噬菌体蛋白 (Yin et al., 2015) 和位点特异性重组酶系统 Cre/loxP 直接克隆基因簇 (Hu et al., 2016)

的重组技术已用于 Photorhabdus 和 Xenorhabdus 共生细菌。此外, 重组酶介导的基因组工程 (Chassis-independent recombinase-assisted genome engineering, CRAGE) 已应用于将共生细菌复杂生物合成基因簇高效整合到细菌的染色体中, 获得高效表达目的蛋白的宿主菌 (Liu et al., 2020)。

4 昆虫病原线虫与共生细菌的专化性

我国在昆虫病原线虫与共生细菌的专化性领域的研究主要集中于如下方面: 共生细菌的信息专化性, 共生细菌的营养专化性, 共生细菌的定殖专化性和共生细菌的杀线虫专化性。

国外学者业已发现, 昆虫血淋巴和共生细菌培养物中含有信号物质 (Food signal) 诱导 IJ 恢复至取食态 (Strauch and Ehlers, 1998; Ciche and Ensign, 2003; Hirao and Ehlers, 2009)。Xenorhabdus 属细菌产生的信号物质不能诱导异小杆线虫发育, 而 Photorhabdus 属中的一些共生菌菌株可诱导 S. carpocapsae 线虫发育 (Ehlers and Han, 1998)。不同种共生细菌产生不同的食物信号, 目前未发现一种共生菌可诱导所有非特异共生的 IJ 恢复 (Han and Ehlers, 1999)。

斯氏属线虫可在高营养的人工培养基中 (Ehlers et al., 1990; Friedman, 1990; Lunau et al., 1993) 和无菌活体昆虫中 (Han and Ehlers, 1999) 无菌繁殖; 但是, 异小杆属线虫对共生细菌具有严格的特异性, 必须在 Photorhabdus 属共生细菌或存在其代谢产物的培养基中才能繁殖 (Akhurst and Boemare, 1990; Ehlers et al., 1990; Han et al., 1990; Aguilera and Smart, 1993; Han et al., 2000)。采用 Tn5 转座方法构建了共生细菌 P. luminescens LN2 突变体库, 筛选出一个对其共生线虫 H. indica LN2 的生长繁殖有显著促进作用的突变菌株, 为构建支持线虫高产的菌株提供了关键技术 (胡素丽等, 2014)。

共生细菌特异性定殖于 IJ 肠道。昆虫病原

线虫都携带特定的共生菌菌株。利用非特异共生细菌可获得肠道未携带细菌的无菌感染期线虫 (Ehlers and Han, 1998; Sicard *et al.*, 2004)。

当两种线虫同时感染一头昆虫寄主时, 两种线虫能否共存或对抗是有趣的问题。当昆虫病原线虫与非特异性共生的共生细菌共同培养时, 一些共生细菌菌株能够产生对非特异性线虫有致死作用的毒素, 比如 *H. indica* LN2 线虫的共生细菌会产生毒素致死 *H. bacteriophora* H06 线虫 (Han and Ehlers, 1999; Qiu *et al.*, 2009)。研究发现, *H. indica* LN2 线虫共生细菌的 *namA* 基因参与了杀线虫毒素的合成 (Qiu *et al.*, 2009)。*P. luminescens* LN2 的 *rpoS* 基因也参与致死 *H. bacteriophora* H06 线虫, 以及 *H. indica* LN2 线虫正常的感染期恢复和发育, 但是, *rpoS* 基因未影响共生细菌在 LN2 和 H06 感染期线虫的定植, 显然 *rpoS* 基因在昆虫病原线虫与共生细菌的互惠关系中发挥了关键作用 (Qiu *et al.*, 2016)。

5 昆虫病原线虫的环境抗性与机制

温湿度、土壤渗透压、紫外线辐射等多种环境因素影响线虫的生存、繁殖及对目标昆虫的感染力 (武海斌等, 2015c; 王杰等, 2021)。我国科研人员在昆虫病原线虫环境耐受品系筛选、昆虫病原线虫响应环境压力的行为和生理生化机制及分子机理方面开展了一系列的研究。

5.1 高温耐受性

杨平和李素春 (1988) 最早研究了温度对 *S. feltiae* 线虫发育、繁殖和感染力的影响, 发现在 10 °C 以下 30 °C 以上不能发育至成虫, 最适生长温度为 25 °C。刘奇志等 (2009) 研究了 *S. longicaudum* BPS 的生物学特性和耐热性。Ma 等 (2013) 比较了 31 个品系的环境耐受性, 发现 *S. carpocapsae*、*S. ceratophorum*、*S. longicaudum*、*H. indica* 和 *H. bacteriophora* 比 *S. feltiae*、*S. hebeiense*、*S. monticolum* 和 *H. megidis* 等品系更耐热。张红玉 (2016) 发现, *H. bacteriophora* JNR 线虫有较好的耐热性, 38 °C 条件下能存活近 32 h

且仍具有侵染活力。Yan 等 (2018) 比较了对黄曲条跳甲幼虫高毒力的 4 种昆虫病原线虫的环境耐受力, 发现采自海南的 *S. pakistanense* 94-1 的耐热能力最强。马娟等 (2022) 发现 *S. tielingense* LFS65、*S. ceratophorum* HQA87 和 *S. carpocapsae* All 线虫对 40 °C 的耐性较强。陈澄等 (2023) 报道 *S. carpocapsae* X-94, 40 °C 下处理 5 h 后存活率为 33%。总体而言, 采自热带地区的昆虫病原线虫品系具有较高的耐热性。在耐热机理方面, 郭笑笑 (2020) 研究发现 *S. ceratophorum* HQA-87 的抗逆性基因 *HSP20* 和 *CL9144* 均与线虫耐高温抗性有关, 但基因沉默结果显示这两个抗逆性基因与线虫致病力不相关。Xie 等 (2020) 研究了经热驯化的 *S. carpocapsae* All 线虫在 38 °C 高温胁迫 5 h 后的转录组, 发现了多个差异表达基因。

5.2 耐低温及抗冻能力

李春杰等 (2011) 在黑龙江省分离到 *H. bacteriophora* HBN 和 *H. bacteriophora* ZT 两个耐寒昆虫病原线虫品系, 发现介质的含水量对 *H. bacteriophora* HBN 冷冻存活率产生显著影响 (李春杰等, 2014), 而适当的预处理可提高 *H. bacteriophora* HBN 和 *H. bacteriophora* NJ 线虫的冷冻存活率 (李春杰等, 2015)。Ma 等 (2013a) 比较发现 *H. megidis*、*H. bacteriophora* 和 *S. carpocapsae* 比测定的其他昆虫病原线虫表现出更好的耐寒性。张红玉 (2016) 研究发现, *H. bacteriophora* JNR 在低温下的侵染力随温度升高而增加。陈桂梅 (2021) 发现, 不同昆虫病原线虫耐寒能力强弱依次为 *S. monticolum* Sm、*S. feltiae* SN、*S. carpocapsae* All、*H. bacteriophora* Hb、*S. glaseri* KG。马娟等 (2022) 发现, 在 -5 °C 处理后, *Steinernema* sp. ZLS3 和 *S. carpocapsae* 线虫的死亡率低于 *S. feltiae* LFS67、*S. feltiae* JY90 和 *H. beicherriana* NCWZ1 等线虫。

张旭霞 (2018) 发现, 阶段性降温驯化后, *S. litorale* 线虫在 -20 °C 的冷冻存活率显著提高, 体内可溶性糖、脂肪和海藻糖含量在驯化后增加, 而蛋白质和糖原含量无显著变化。陈桂梅

(2021) 分析了 *H. bacteriophora* Hb、*S. carpocapsae* All、*S. monticolum* Sm、*S. feltiae* SN 的水通道蛋白基因在冷冻胁迫下的不同转录响应。

5.3 紫外线辐射耐性及保护剂

紫外线是昆虫病原线虫防治地上害虫时最重要的影响因子。钱秀娟等 (2019a) 研究紫外线辐射对昆虫病原线虫存活率和致病力的影响, 发现昆虫病原线虫不同品系均对紫外辐射具有一定的耐受能力。肖建飞等 (2021) 发现木质素磺酸钙能显著提高 *S. litorale* 抗紫外胁迫的能力。Wu 等 (2023a) 发现氨基化二氧化钛 (Titania-NH₂) 纳米粒子可保护 *S. carpocapsae* 免受紫外线辐射。Ha 等 (2025) 发现 0.2% 葡萄籽提取物与 0.1% 阻燃凝胶的组合对 *S. carpocapsae* 露于 3 种波长的人工紫外线辐射 (UVA、UVB、UVC) 下有最好的保护效果。丁倩 (2024) 发现农用植物油可提高 *H. bacteriophora* Hb、*H. beicherriana* H06、*H. indica* LN2、*S. carpocapsae* All 和 *S. feltiae* SN 线虫在紫外线辐射处理下的存活率。

5.4 耐受脱水能力及机制

不同种的昆虫病原线虫, 以及同种不同品系的昆虫病原线虫, 耐受脱水的能力不同 (冯世鹏和韩日畴, 2005; 颜珣和韩日畴, 2010)。陈松笔等 (1999) 筛选获得抗干燥能力强的线虫种类。邢玉芳等 (2009) 测定了甘肃 8 种昆虫病原线虫的抗干燥能力。李宁等 (2011) 测定了 10 个昆虫病原线虫品系的抗干燥能力, *S. carpocapsae* GA 品系的存活率最高, *Heterorhabditis* sp. CZ 和 XH-9 品系则全部死亡。Ma 等 (2013) 发现, *S. carpocapsae*、*S. hebeiense* 和 *S. ceratophorum* 具有良好的耐脱水性。吴霞等 (2019) 报道, 异小杆线虫抗低湿胁迫能力弱于斯氏线虫, 斯氏线虫 *S. kraussei* 0657L 抗低湿胁迫能力强于测定的其他线虫, 低湿胁迫对 *S. feltiae* 0619HT 和 *H. megidis* 0627M 的繁殖无显著影响。

不同昆虫病原线虫品种之间渗透压耐性存在差异, *S. carpocapsae* All 和 *S. scapterisci* SS

最耐受渗透压溶液处理 (Yan *et al.*, 2010, 2011)。马娟等 (2022) 测定不同地区的昆虫病原线虫的抗干燥能力, 发现 *Steinernema* sp. ZLS3、*S. carpocapsae* All 和 *S. ceratophorum* HQA87 在 25% 甘油溶液中的存活率高于测定的其他线虫。

昆虫病原线虫脱水胁迫下进入休眠状态后, 体表和体内发生了显著变化 (陈松笔等, 2001b), 与耗氧量相关 (陈松笔等, 2001a)。不同昆虫病原线虫应对低湿胁迫的避逆策略不同, 在 30% 低湿环境中, *S. kraussei* 0657L 表现为干燥逃避, *H. brevicaudis* 0641TY 则表现为低湿休眠 (张晓波等, 2023)。 *S. carpocapsae* BJ 线虫在高渗溶液中脱水后进入休眠状态, 其酯酶活性降低, 总糖和糖原含量均产生改变 (陈松笔等, 2000)。甘油处理后脱水的 *S. carpocapsae* Agriotos 合成甘油和海藻糖等保护化合物 (Qiu *et al.*, 2000; Qiu and Bedding, 2002)。吴霞 (2023) 也发现, 低湿胁迫会引起 *S. kraussei* 0657L 和 *H. brevicaudis* 0641TY 体内总脂肪、海藻糖和蛋白质的积累, 同时能量物质与存活率和致病力之间存在着一定的相关性。

在分子响应机制方面, *S. feltiae* IS-6 线虫可通过 24% 甘油溶液处理进入渗透脱水状态, UNC-87、肌动蛋白 (Actin)、蛋白酶体调节颗粒 (ATP 酶样) [Proteasome regulatory particle (ATPase-like)]、GroEL 伴侣蛋白 (GroEL chaperonin)、GroES 共伴侣蛋白 (GroES co-chaperonin) 以及转座酶家族成员 (Transposase family member) 等蛋白的上调表达与渗透压胁迫的响应相关 (Chen *et al.*, 2005, 2006b)。 *S. carpocapsae* 的 *lea* 和 *sod-2* 基因在耐热性较弱的 MR7 品系中的表达量显著高于 All 品系, 而 *t19d7* 的表达量则显著降低 (Yan *et al.*, 2011)。对渗透压耐性差异显著的 4 种昆虫病原线虫 *S. carpocapsae*、*S. loci*、*H. bacteriophora* 和 *H. indica* 中 *fat-2* 的表达情况分析结果显示, 该基因在渗透压耐性最强的 *S. carpocapsae* 线虫中表达量最高, 可能与渗透压耐性差异直接相关 (Yan *et al.*, 2012)。吴霞 (2023) 报道 *TRE* 和 *CAT* 基因能

够调控 *S. kraussei* 0657L 体内能量物质以响应低湿胁迫。黄山 (2023) 通过 RNA 干扰 (RNA interference, RNAi) 技术验证了 *CL2037* 和 *CL2161* 基因与 *S. ceratophorum* HQA-87 线虫的干燥抗性有关。研究表明, 昆虫病原线虫对脱水响应的分子机制比较复杂, 通过基因转化和添加单个基因来改善昆虫病原线虫的脱水耐受性不太可能实现。

6 昆虫病原线虫的应用

昆虫病原线虫种类会影响其对不同害虫的毒力效果, 应用昆虫病原线虫时应针对特定靶标害虫筛选高毒力的品种和品系, 以实现靶标害虫的有效控制。我国学者针对我国农业中的一系列地下害虫、钻蛀害虫以及食叶害虫, 筛选获得相应的高毒力昆虫病原线虫 (表 1)。

表 1 昆虫病原线虫防控的主要害虫
Table 1 Major pests controlled by entomopathogenic nematodes

目标昆虫 Target insect	线虫种或品系 Nematode species/strain	参考文献 Reference
桃小食心虫 <i>Carposina nipponensis</i>	<i>H. bacteriophora</i> H06	李爱华等, 2012
梨小食心虫 <i>Grapholita molesta</i>	<i>S. carpocapsae</i> All	严凯等, 2017
韭菜迟眼蕈蚊 <i>Bradysia odoriphaga</i>	<i>H. bacteriophora</i> ZT, <i>S. carpocapsae</i> , <i>S. feltiae</i> SN, <i>H. indica</i> LN2	李春杰等, 2013; Ma <i>et al.</i> , 2013b; 潘凤娟等, 2013; 张燕等, 2014; Bai <i>et al.</i> , 2016; Yan <i>et al.</i> , 2019
华北大黑鳃金龟 <i>Holotrichia oblita</i>	<i>H. beicherriana</i> LF	李而涛等, 2018
铜绿丽金龟 <i>Anomala corpulenta</i>	<i>H. beicherriana</i> D26-6, <i>S. carpocapsae</i>	张维伟和赵福永, 2021; 白勇章等, 2024
暗黑鳃金龟 <i>Holotrichia parallela</i>	<i>H. bacteriophora</i> D26-6, <i>H. beicherriana</i>	李而涛等, 2020; 张彩铃等, 2024
小云斑鳃金龟 <i>Polyphylla gracilicornis</i>	<i>S. longicaudum</i> X-7	范芳芳等, 2015
白星花金龟 <i>Postosia brevitarsis</i>	<i>S. carpocapsae</i> All	李长青等, 2011
小地老虎 <i>Agrotis ipsilon</i>	<i>S. carpocapsae</i> , <i>H. indica</i> LN2	Yan <i>et al.</i> , 2014; 武海斌等, 2015b; 李根等, 2017; 谷星慧等, 2021
黄地老虎 <i>Agrotis segetum</i>	<i>S. glaseri</i> ib	张安邦等, 2015
八字地老虎 <i>Xestia cnigrum</i>	<i>S. carpocapsae</i> -2	张思佳等, 2013
草地贪夜蛾 <i>Spodoptera frugiperda</i>	<i>S. carpocapsae</i> All, <i>S. feltiae</i> , <i>H. indica</i> , <i>H. megidis</i> 0872L	梁铭荣等, 2020; 杨淋凯等, 2021; 蔡佳仪等, 2023; Guo <i>et al.</i> , 2023
斜纹夜蛾 <i>Spodoptera litura</i>	<i>S. longicaudum</i> X-7, <i>H. indica</i> 212-2	Yan <i>et al.</i> , 2020; Sun <i>et al.</i> , 2021
甜菜夜蛾 <i>Spodoptera exigua</i>	<i>H. indica</i> LN2, <i>S. carpocapsae</i>	刘敏等, 2025
二点委夜蛾 <i>Athetis lepigone</i>	<i>H. bacteriophora</i> HB8	雷利平等, 2013
番茄潜麦蛾 <i>Tuta absoluta</i>	<i>S. carpocapsae</i> All	郭文秀等, 2024
舞毒蛾 <i>Lymantria dispar</i>	<i>S. carpocapsae</i> All	滑莎等, 2018
腰果细蛾 <i>Acrocercops syngamma</i>	<i>S. carpocapsae</i> All	张中润等, 2016
苹果蠹蛾 <i>Cydia pomonella</i>	<i>S. riobrave</i> , <i>S. carpocapsae</i>	张金花等, 2024
小线角木蠹蛾 <i>Holcocerus hippophaecolus</i>	<i>S. carpocapsae</i>	路常宽等, 2004
二化螟 <i>Chilo suppressalis</i>	<i>S. carpocapsae</i> N-Yz1, <i>S. carpocapsae</i> XC-4	刘琴等, 2024; 刘洋等, 2025
腰果云翅斑螟 <i>Hyalospila leuconeurella</i>	<i>S. carpocapsae</i> All	张中润等, 2017
草地螟 <i>Loxostege sticticalis</i>	<i>S. carpocapsae</i> All	李朔涵, 2023

续表 1 (Table 1 continued)

目标昆虫 Target insect	线虫种或品系 Nematode species/strain	参考文献 Reference
草原毛虫 <i>Gynaephora alpherakii</i>	<i>S. feltiae</i> SN, <i>S. carpocapsae</i> All	白海涛等, 2020
黄曲条跳甲 <i>Phyllotreta striolata</i>	<i>S. carpocapsae</i> All	杨凤丽等, 2020; 赵丽稳等, 2024
红棕象甲 <i>Rhynchophorus ferrugineus</i>	<i>S. carpocapsae</i> All, <i>H. bacteriophora</i> H06	钟宝珠等, 2020; Lv <i>et al.</i> , 2025
茶籽象甲 <i>Curculio chinensis</i>	<i>S. carpocapsae</i> All	王轶等, 2019
小蜂巢甲虫 <i>Aethina tumida</i>	<i>S. carpocapsae</i> All	韩文素等, 2021
黄粉虫 <i>Tenebrio molitor</i>	<i>S. litorale</i>	詹发强等, 2020
曲牙土天牛 <i>Dorysthenes hydropicus</i>	<i>H. bacteriophora</i> H06	徐常青等, 2010
松褐天牛 <i>Monochamus alternatus</i>	<i>S. carpocapsae</i> All	张彦龙等, 2012
核桃云斑天牛 <i>Batocera lineolata</i>	<i>S. carpocapsae</i>	陈贤群等, 2016
甘薯蚁象 <i>Cylas formicarius</i>	<i>S. feltiae</i>	于海滨等, 2012
臭椿沟眶象 <i>Eucryptorrhynchus brandti</i>	<i>S. carpocapsae</i> All	郑惠中, 2021
白纹伊蚊 <i>Anopheles albopictus</i>	<i>S. abbasi</i>	Liu <i>et al.</i> , 2020
红火蚁 <i>Solenopsis invicta</i>	<i>H. bacteriophora</i> , <i>S. riobrave</i> , <i>S. carpocapsae</i> All	Wu <i>et al.</i> , 2023b; Li <i>et al.</i> , 2024c; 易松望等, 2024
南瓜实蝇 <i>Bactrocera tau</i>	<i>S. feltiae</i>	Liu <i>et al.</i> , 2025

根据王杰等 (2021) 改进。
Improved according to Wang *et al.*, 2021.

6.1 昆虫病原线虫与化学药剂共用

在害虫防治实践中, 昆虫病原线虫可与兼容的化学药剂混用, 以扩大防治谱或提高防治效果。我国应用昆虫病原线虫与低毒化学药剂共同防治害虫, 取得较好的效果 (表 2)。

6.2 昆虫病原线虫与其它微生物混用

在我国, 开展了将昆虫病原线虫与其他昆虫病原如苏云金芽孢杆菌 *Bacillus thuringiensis* (Hatting *et al.*, 2019)、球孢白僵菌 *Beauveria bassiana* (张伟等, 2017)、绿僵菌 *Metarhizium anisopliae* (黄亚川等, 2020) 等协同防控害虫的研究。苏云金芽孢杆菌产生的毒素引起黏虫 *Mythimna separata* 中肠组织病变, 降低黏虫的免疫力, 有利于昆虫病原线虫的感染, 因此两者联用对黏虫具有协同防控效果 (李而涛等, 2019)。PI3K/Akt/CncC 信号通路介导昆虫病原线虫和苏云金芽孢杆菌对暗黑鳃金龟 *Holotrichia parallela* 幼虫的感染 (Li *et al.*, 2023)。此外还发现,

腐生真菌腐皮镰刀菌 *Fusarium solani* 可吸引并促进 *S. diaprepesi* 线虫感染寄主昆虫 (Wu *et al.*, 2018)。

6.3 昆虫病原线虫田间应用的典型案例

在我国, 已利用昆虫病原线虫防治桃小食心虫 (杨平等, 2000)、黄曲条跳甲 (Yan *et al.*, 2013)、韭菜迟眼蕈蚊 *Bradysia odoriphaga* (Ma *et al.*, 2013b)、小地老虎 *Agrotis ipsilon* (Yan *et al.*, 2014)、华北大黑鳃金龟 *Holotrichia oblita* (Guo *et al.*, 2017)、草地贪夜蛾 (唐璞等, 2019) 等多种害虫。典型的成功案例包括利用昆虫病原线虫田间防控桃小食心虫、蛴螬和韭蛆。

6.3.1 桃小食心虫 桃小食心虫主要危害苹果、桃、杏、李、山楂、枣、石榴等, 已成为我国苹果生产中发生面积最大、危害最严重的害虫之一 (孙丽娜等, 2018)。早在 20 世纪 80 年代, 国内便应用昆虫病原线虫大面积防治桃小食心虫, 防控效果显著。在山东泰安、寿光等地 5 个果园苹果生产中应用 *S. carpocapsae* Agriotes 线虫在

表 2 昆虫病原线虫与化学药剂联用防控的主要害虫
Table 2 Combination of entomopathogenic nematodes and chemicals for the control of insect pests

目标昆虫 Target insect	线虫种和品系 EPN species/strain	化学药剂种类 Chemical insecticides	参考文献 Reference
韭菜迟眼蕈蚊 <i>Bradysia odoriphaga</i>	<i>S. feltiae</i> SN	噻虫嗪 Thiamethoxam	武海斌等, 2018
	<i>S. feltiae</i> SN	吡虫啉 Imidacloprid	Yan <i>et al.</i> , 2019; 付在秋等, 2020
	<i>S. feltiae</i> 0619HT	吡虫啉 Imidacloprid	钱秀娟等, 2019b
	<i>S. carpocapsae</i> All	溴氰虫酰胺 Cyantraniliprole	翟婧等, 2020
卵圆齿爪鳃金龟 <i>Holotrichia ovata</i>	<i>S. longicaudum</i> X-7	除虫脲 Diflubenzuron	张中润等, 2006
华北大黑鳃金龟 <i>Holotrichia oblita</i>	<i>H. bacteriophora</i>	吡虫啉 Imidacloprid	Guo <i>et al.</i> , 2017c
褐纹甘蔗象 <i>Rhabdoscelus lineaticollis</i>	<i>S. carpocapsae</i> All	吡虫啉 Imidacloprid	王果红等, 2007
黄曲条跳甲 <i>Phyllotreta striolata</i>	<i>S. carpocapsae</i> All	溴氰虫酰胺 Cyantraniliprole	金立新等, 2019
黄粉虫 <i>Tenebrio molitor</i>	<i>S. carpocapsae</i> All	茚虫威 Indoxacarb	陈瑞等, 2019
红火蚁 <i>Solenopsis invicta</i>	<i>S. riobrave</i>	联苯菊酯与噻虫胺混合物 Mixture of bifenthrin and clothianidin	Wu <i>et al.</i> , 2024
臭椿沟眶象 <i>Eucryptorrhynchus brandti</i>	<i>S. longicaudum</i> X-7	氯氰菊酯 Cypermethrin, 吡虫啉 Imidacloprid	郑惠中, 2021
烟粉虱 <i>Bemisia tabaci</i>	<i>S. carpocapsae</i> All	柠檬烯可溶液剂 Limonene solution	陈向荣等, 2021

根据王杰等 (2021) 改进。
Improved according to Wang *et al.*, 2021.

1.0 亿/667 m² 剂量下对桃小食心虫的控制效果优于化学防控区 (刘杰, 1994; 杨平等, 2000)。在河北地区, *S. feltiae* Beijing 线虫在 1.0 亿/667 m² 剂量下能够有效降低虫果率, 防控效果优于化学药剂 (刘新生等, 1993)。刘峥等 (1998) 分离到一种对桃小食心虫高毒力的本土昆虫病原线虫 *Steinernema* sp., 在河南郑州、新乡、博爱等地开展 3 年的田间防控试验也表明, 该线虫较常规化学药剂能够减少 22%-68% 桃小食心虫的出土数量及 39%-76% 的虫果数量。桃小越冬幼虫出土一般在 5 月末 6 月初, 此时土壤温度和含水量均适于线虫的存活及侵染。目前, 苹果产业正经历由传统果园向现代果园发展历程, 桃小食心虫的防治已转套袋防治, 昆虫病原线虫在现代果园中防治桃小食心虫还需进一步加强研究。

6.3.2 蛴螬 蛴螬是昆虫病原线虫最早应用及应用研究最为广泛的目标地下害虫 (Glaser and Fox, 1930), 一系列种类或品系如 *S. glaseri* (Converse and Grewal, 1998)、*S. scarabaei*

(Cappaert and Koppenhöfer, 2003; Koppenhöfer *et al.*, 2006)、不同品系的 *H. bacteriophora* (Koppenhöfer *et al.*, 2000, 2002; Koppenhöfer and Fuzy, 2003; Grewal *et al.*, 2004)、*H. zealandica* (Grewal *et al.*, 2004) 等已被成功地应用于蛴螬的防治。在我国, 一些本土昆虫病原线虫对蛴螬有很好的控制效果。花生田施用 *S. longicaudum* X-7、*H. bacteriophora* H06 IJ 线虫 (0.5-1.0 亿/667 m²) 能够有效控制暗黑鳃金龟、华北大黑鳃金龟 *H. oblita*, 使蛴螬减退率达 88% 以上 (Guo *et al.*, 2013, 2015)。烟草田中应用 *S. longicaudum* X-7 能够有效减少蛴螬虫口密度, 保苗效果显著 (李根, 2017)。金银花田施用 *S. longicaudum* X-7 能够持续控制蛴螬危害, 植株死亡率显著降低 (Li *et al.*, 2023)。一些环境友好型药剂如氯化烟酰胺类 (王玉东等, 2012)、昆虫生长调节剂类药剂除虫脲 (张中润等, 2006) 与线虫混用具有协同增效作用, 可提高线虫对蛴螬的防治效果。田间施用昆虫病原线虫的时间节

点非常关键,应根据当地作物蛴螬的发生危害规律,选择 1-2 龄幼虫期,采用滴灌装备将线虫悬液沿植株根茎顺流入根系土壤中发挥持续性防控作用 (Guo *et al.*, 2015)。

6.3.3 韭蛆 国内学者开展了大量应用昆虫病原线虫防治韭蛆 (迟眼蕈蚊幼虫) 的研究工作,筛选到了多种对韭蛆高毒力的线虫种类,如 *S. feltiae* 多个品系 (杨秀芬等, 2004; Ma *et al.*, 2013)、*H. indica* LN2 (孙瑞红等, 2004; Ma *et al.*, 2013b)、*H. bacteriophora* ZT (李春杰等, 2013) 等。一系列田间应用技术及配套技术的持续研究,支撑了昆虫病原线虫对韭蛆的田间成功控制。杨秀芬等 (2004) 对 *S. feltiae* PS4 进行田间小区试验测试,对韭蛆防治效果为 55.8%。Ma 等 (2013) 针对昆虫病原线虫防治韭蛆开展了系统研究,发现 *S. feltiae* JY-17 线虫对韭蛆的致死率可达 97%。赵国玉 (2013) 对露天栽培韭菜田中夏季应用 *H. indica* LN2 线虫对韭蛆的持续性控制效果进行了研究,发现 *H. indica* LN2 IJ 线虫按照 1.5 亿/667 m² 施用 14 d 后韭蛆虫口密度减少 50%,与辛硫磷无显著差异,同时,线虫对韭蛆具有持续性的防治效果,直至初秋韭蛆虫口密度始终保持在 100 头/m² 以下的较低密度。对于保护地栽培的韭菜生产,在春、秋、冬三个关键节点,施用添加 1/10 田间推荐剂量吡虫啉的 *S. feltiae* SN IJ 线虫 2.0 亿/667 m² 表现出优于灭幼脲的防控效果,能够实现韭蛆的周年有效控制 (赵国玉等, 2013; Yan *et al.*, 2019)。

7 昆虫病原线虫活力与环境因子的关系

7.1 昆虫病原线虫与昆虫寄主

H. bacteriophora 线虫侵染导致暗黑鳃金龟和大黑鳃金龟幼虫脂肪体和中肠细胞出现明显病变 (吴文丹等, 2015d)。暗黑鳃金龟的免疫相关基因 *PGRP-SC1*、*PPAE-I* 和 *GST* 参与了对 *H. beicherriana* 线虫感染的抵抗 (Li *et al.*, 2021b)。昆虫病原线虫可抑制水椰八角铁甲 *Octodonta nipae* 酚氧化酶系统的激活 (Sanda *et*

al., 2018; Sanda and Hou, 2023), 以及草地贪夜蛾 (李而涛等, 2022) 和冷杉梢斑螟 *Dioryctria abietella* (聂振业等, 2023) 的多酚氧化酶、超氧化物歧化酶、过氧化氢酶、乙酰胆碱酯酶、谷胱甘肽 S-转移酶等的变化。*S. feltiae* 线虫感染大蜡螟幼虫后,昆虫的色氨酸 (Trp) 途径被显著激活,血淋巴中的犬尿氨酸 (Kyn) 和 3-羟基苯甲酸 (3-HAA) 含量显著升高; 3-HAA 对大蜡螟幼虫高毒性 (Zhang *et al.*, 2022)。*S. carpocapsae* 侵染可导致榛实象甲 *Curculio dieckmanni* 幼虫相关基因和代谢物的改变 (Wang *et al.*, 2025)。*H. megidis* 比 *S. feltiae* 对大蜡螟幼虫表现出更强的趋性,且寄主的存在对线虫的迁移能力有影响 (吴霞等, 2021)。

7.2 昆虫病原线虫与活性化合物

蛴螬类线虫信息化合物具有调控线虫聚集、趋避、交配、耐受态幼虫形成及扩散传播等行为的功能 (Jeong *et al.*, 2005; Kim *et al.*, 2009; Kaplan *et al.*, 2012; Hartley *et al.*, 2019; Oliveira-Hofman *et al.*, 2019)。研究发现,某些蛴螬影响 IJ 的恢复、扩散和在琼脂平板和液体培养基中的产量 (Kong *et al.*, 2022; Wang *et al.*, 2022)。此外,蛴螬和共生细菌调控大蜡螟幼虫的免疫响应 (Chantab *et al.*, 2024)。

研究已经发现,异戊醇对多种昆虫病原线虫具有排斥作用 (Baiocchi *et al.*, 2020), 而寄主昆虫本身分泌的化学物质可以吸引昆虫病原线虫。红棕象甲 *Rhynchophorus ferrugineus* 利用分泌的化学信号吸引 *S. carpocapsae* 线虫 (Santhi *et al.*, 2021)。线虫感染致死的玉米根叶甲 *Diabrotica virgifera virgifera* 虫尸的挥发性物质对其他玉米根叶甲幼虫具有吸引作用,有利于线虫感染更多的寄主并成功繁殖 (Zhang *et al.*, 2019c; Santhi *et al.*, 2021)。 α -松油醇影响群体免疫 (Social immunity), 增加昆虫病原线虫对白蚁的致病性 (Zeng *et al.*, 2023)。甘油磷酸胆碱 (Glycerophosphocholine) 参与了 *S. kraussei* 0657L 的寿命调控,并通过激活胰岛素/IGF-1 信号通路和下游 PI3K/MAPK 激酶在抵抗外部应激中发挥重要作用,为提高 *S. kraussei* 的商业功效

提供了新的思路 (Li *et al.*, 2024d)。被昆虫病原线虫感染的昆虫尸体所释放的挥发性化合物影响了南方根结线虫的发育 (Li *et al.*, 2024a)。由于多重毒性作用, *S. feltiae* 线虫摄入聚苯乙烯微塑料后会损害其适合度 (Li *et al.*, 2024b)。绿原酸被认为可能通过胰岛素/IGF-1 信号通路提高 *S. kraussei* 的抗氧化能力并延长其寿命 (陈红等, 2024)。

7.3 昆虫病原线虫与环境非生物因素

田间应用昆虫病原线虫时, 土壤的结构和质地、湿度、温度、田间施用剂量以及施用方法等因素直接影响昆虫病原线虫的应用效果 (Půža and Mráček, 2005; 武海斌等, 2015c)。 *S. feltiae* SN 在 15-20 °C 的温度范围内, 比 *H. indica* LN2 对韭菜迟眼蕈蚊感染效果更强, 表明两种昆虫病原线虫的最佳侵染温度有差异 (Yan *et al.*, 2019)。 *Steinernema* 线虫在 25 °C、相对湿度为 50.5% 环境中处理 9 h 后存活率仍较高, 而 *Heterorhabditis* 线虫几乎没有存活 (邢玉芳等, 2009)。5 龄二点委夜蛾 *Athetis lepigone* 幼虫在 15% 土壤含水量时对 *H. bacteriophora* HB8 线虫最敏感 (雷利平等, 2013), 当土壤含水量为 20% 时 *S. glaseri* 线虫对黄地老虎 *A. segetum* 幼虫的防治效果最好 (张安邦, 2015)。可通过添加抗干燥剂及抗蒸发剂以提高昆虫病原线虫防治害虫的效果 (金永玲和韩日畴, 2009; 金永玲等, 2013)。土壤质地影响昆虫病原线虫的分布、扩散和对昆虫的感染毒力 (王庆雷等, 1995; 钱秀娟等, 2005; 陈书龙等, 2006)。沙壤土和壤土更适合昆虫病原线虫建立种群 (钱秀娟等, 2014)。

8 展望

作为新型的昆虫天敌类生物农药, 昆虫病原线虫及其共生细菌已经商业生产, 应用于全球多种有害生物的防控。展望未来, 这类制剂在我国的应用仍需要高度关注如下趋势:

1) 政策指引: 生物农药的广泛应用离不开环境保护政策的支持。“绿水青山就是金山银山”

发展理念已写进党章, 在这一核心理论的指导下, 我国生物农药包括昆虫病原线虫的发展前景广阔。

2) 技术优化和成本控制: 昆虫病原线虫保持持续商业竞争力的关键因素是基于昆虫病原线虫、共生细菌以及寄主昆虫的交互作用深入研究基础上, 更优化的培养系统且更高效的贮存技术; 同时, 通过资源筛选、杂交、遗传改造等方法获得更耐受干燥、极端温度、紫外线等环境条件的昆虫病原线虫品系。

3) 市场信誉: 像其它生物农药一样, 昆虫病原线虫的控制对象是害虫、植物病原等, 评价其价值的关键是被保护农产品的质量和价格。消费者愿意为了健康高消费被生物农药保护的农产品。但市场上不应该出现假冒被生物农药保护的农产品, 这样会动摇消费者的信心, 从而摧毁被生物农药保护的农产品的信誉。这也是未来尤其需要关注的重点。

参考文献 (References)

- Aguillera MM, Smart GC, 1993. Development, reproduction and pathogenicity of *Steinernema scapterisci* in monoxenic culture with different species of bacteria. *Journal of Invertebrate Pathology*, 62(3): 289–294.
- Ahuja A, Kushwah J, Mathur C, Chauhan K, Dutta TK, Somvanshi VS, 2021. Identification of galtoxin, a new protein toxin from *Photorhabdus* bacterial symbionts of *Heterorhabditis* nematodes. *Toxicon*, 194: 53–62.
- Akhurst RJ, 1980. Morphological and functional dimorphism in *Xenorhabdus* spp., bacteria symbiotically associated with the insect pathogenic nematodes *Neosapientia* and *Heterorhabditis*. *Microbiology*, 121(2): 303–309.
- Akhurst RJ, Boemare NE, 1990. Biology and Taxonomy of *Xenorhabdus*, Entomopathogenic Nematodes in Biological Control. Boca Raton: Chemical Rubber Company Press. 75–90.
- Anes KM, Patil J, Babu M, Aarthi N, Gotyal BS, Josephraj Kumar A, Mhatre PH, Sajan JV, Gowda MT, Půža V, 2025. *Steinernema keralense* n. sp., a new entomopathogenic nematode (Nematoda: Steinernematidae) from India. *Journal of Helminthology*, 99: e31.
- Bai GY, Xu H, Fu YQ, Wang XY, Shen GS, Ma HK, Feng XL, Pan J, Gu XS, Guo YZ, Ruan WB, Shapiro-Ilan DI, 2016. A comparison of novel entomopathogenic nematode application methods for

- control of the chive gnat, *Bradysia odoriphaga* (Diptera: Sciaridae). *Journal of Economic Entomology*, 109(5): 2006–2013.
- Bai HT, Wang W, Wei XJ, Han RC, Xu CT, 2020. Pathogenicity of entomopathogenic nematodes against *Gynaephora alpherakii* larvae by bioassay. *Journal of Environmental Entomology*, 42(5): 1269–1274. [白海涛, 王伟, 魏希杰, 韩日畴, 徐成体, 2020. 昆虫病原线虫对草原毛虫幼虫的致病力. 环境昆虫学报, 42(5): 1269–1274.]
- Bai YZ, Wang WJ, Wang ZM, Li SH, Li KB, Cao YZ, Wang SS, Song LW, Yin J, 2024. Screening of entomopathogenic nematodes with high virulence to *Anomala corpulenta*. *Plant Protection*, 50(1): 334–338, 355. [白勇章, 王文杰, 王志敏, 李朔涵, 李克斌, 曹雅忠, 王森山, 宋丽雯, 尹姣, 2024. 对铜绿丽金龟具有高致病力的昆虫病原线虫的筛选. 植物保护, 50(1): 334–338, 355.]
- Baiocchi T, Li CJ, Dillman AR, 2020. EPNs exhibit repulsion to prenil in pluronic gel assays. *Insects*, 11(8): 457.
- Baiome BA, Ye XF, Yuan ZY, Gaafar YZA, Melak S, Cao H, 2022. Identification of volatile organic compounds produced by *Xenorhabdus indica* strain AB and investigation of their antifungal activities. *Applied and Environmental Microbiology*, 88(13): e0015522.
- Baniya A, Subkrasae C, Ardpairin J, Anesko K, Vitta A, Dillman AR, 2024. *Steinernema adamsi* n. sp. (Rhabditida: Steinernematidae), a new entomopathogenic nematode from Thailand. *The Journal of Parasitology*, 110(1): 22–39.
- Bedding RA, 1981. Low cost *in vitro* mass production of *Neoaplectana* and *Heterorhabditis* species (Nematoda) for field control of insect pests. *Nematologica*, 27(1): 109–114.
- Bhat AH, Chaubey AK, Askary TH, 2020. Global distribution of entomopathogenic nematodes, *Steinernema* and *Heterorhabditis*. *Egyptian Journal of Biological Pest Control*, 30(1): 31.
- Bhat AH, Machado RAR, Abolafia J, Askary TH, Půža V, Ruiz-Cuenca AN, Rana A, Sayed S, Al-Shuraym LA, 2023a. Multigene sequence-based and phenotypic characterization reveals the occurrence of a novel entomopathogenic nematode species, *Steinernema anantnagense* n. sp. *Journal of Nematology*, 55(1): 20230029.
- Bhat AH, Machado RAR, Abolafia J, Ruiz-Cuenca AN, Askary TH, Ameen F, Dass WM, 2023b. Taxonomic and molecular characterization of a new entomopathogenic nematode species, *Heterorhabditis casmirica* n. sp., and whole genome sequencing of its associated bacterial symbiont. *Parasites & Vectors*, 16(1): 383.
- Bi YH, Gao CZ, Yu ZG, 2018. Rhabdopeptides from *Xenorhabdus budapestensis* SN84 and their nematocidal activities against *Meloidogyne incognita*. *Journal of Agricultural and Food Chemistry*, 66(15): 3833–3839.
- Bogdanovic X, Schneider S, Levanova N, Wirth C, Trillhaase C, Steinemann M, Hunte C, Aktories K, Jank T, 2019. A cysteine protease-like domain enhances the cytotoxic effects of the *Photorhabdus asymbiotica* toxin PaTox. *The Journal of Biological Chemistry*, 294(3): 1035–1044.
- Bowen D, Blackburn M, Rocheleau T, Grutzmacher C, ffrench-Constant RH, 2000. Secreted proteases from *Photorhabdus luminescens*: Separation of the extracellular proteases from the insecticidal Tc toxin complexes. *Insect Biochemistry and Molecular Biology*, 30(1): 69–70.
- Buka-Ouerna, Zhan FQ, Long XQ, Luo M, Hou M, Yang R, 2015. Mass production technology for *Heterorhabditis* NLK-1. *Xinjiang Agricultural Sciences*, 52(9): 1693–1699. [布卡·欧尔娜, 詹发强, 龙宣杞, 罗明, 侯敏, 杨蓉, 2015. 异小杆线虫 NLK-1 大量增殖技术研究. 新疆农业科学, 52(9): 1693–1699.]
- Cai JY, Zhang XH, Liu QZ, Li XY, Tang ZL, 2023. Larval mortality and field trial effect caused by entomopathogenic nematodes on *Spodoptera frugiperda*. *Journal of Biosafety*, 32(1): 56–62. [蔡佳仪, 张小涵, 刘奇志, 李星月, 唐照磊, 2023. 昆虫病原线虫对草地贪夜蛾幼虫的致死率及田间防治效果. 生物安全学报, 32(1): 56–62.]
- Cao LQ, Zhan FQ, Gao YJ, Hou XQ, Bao HF, Hou M, Yang R, Wang N, Long XQ, 2021. The inhibitory effect of *Xenorhabdus nematophilus* on *Botrytis cinerea*. *Microbiology China*, 48(11): 4123–4133. [曹林青, 詹发强, 高宇洁, 侯新强, 包慧芳, 侯敏, 杨蓉, 王宁, 龙宣杞, 2021. 嗜线虫致病杆菌抑制灰葡萄孢的效应. 微生物学通报, 48(11): 4123–4133.]
- Cappaert DL, Koppenhöfer AM, 2003. *Steinernema scarabaei*, an entomopathogenic nematode for control of the European chafer. *Biological Control*, 28(3): 379–386.
- Chang SH, Yang HW, 1998. The effects of nutrition on reproduction ability and nematode yield in continued culture. *Chinese Journal of Biological Control*, 14(3): 105–106. [常韶华, 杨怀文, 1998. 连续传代培养中营养对昆虫病原线虫的影响. 中国生物防治学报, 14(3): 105–106.]
- Chantab K, Rao ZC, Zheng XH, Han RC, Cao L, 2024. Ascarosides and symbiotic bacteria of entomopathogenic nematodes regulate host immune response in *Galleria mellonella* larvae. *Insects*, 15(7): 514.
- Chen C, Hou XQ, Zhan FQ, Yang R, Bao HF, Wang N, Shi YW, Long XQ, 2023. Study on abiotic stress tolerance of three entomopathogenic nematodes and its bacteriostasis of symbiotic bacteria. *Xinjiang Agricultural Sciences*, 60(12): 3072–3079. [陈

- 澄, 侯新强, 詹发强, 杨蓉, 包慧芳, 王宁, 史应武, 龙宣杞, 2023. 三种昆虫病原线虫非生物胁迫耐受特性及其共生菌抑菌作用. 新疆农业科学, 60(12): 3072–3079.]
- Chen GM, 2021. Response analysis of aquaporins in entomopathogenic nematodes under freezing stress. Master dissertation. Guangzhou: South China Agricultural University. [陈桂梅, 2021. 昆虫病原线虫冷冻胁迫下水通道蛋白的响应分析. 硕士学位论文. 广州: 华南农业大学.]
- Chen H, Li XT, Wang XD, Qian XJ, 2024. Regulation of longevity and antioxidant activity in entomopathogenic nematodes *Steinernema kraussei* by chlorogenic acid. *Chinese Journal of Biological Control*, 40(1): 108–116. [陈红, 李茜童, 王兴铎, 钱秀娟, 2024. 绿原酸对昆虫病原线虫卡森氏线虫寿命及抗氧化活性的调控. 中国生物防治学报, 40(1): 108–116.]
- Chen JH, Han RC, 2014. A spraying device for entomopathogenic nematodes. Chinese Invention Patent, ZL201210437336.5. 2014-04-09. [陈镜华, 韩日畴, 2014. 一种昆虫病原线虫喷施设备. 中国发明专利, ZL201210437336.5. 2014-04-09.]
- Chen R, Sheng XR, Wang DZ, Wang YX, Sun WM, Kong LW, Xia B, Wang GD, 2020. Culture bottle for entomopathogenic nematodes. Chinese Utility Model Patent, ZL201921993353.0. 2020-10-09. [陈瑞, 盛锡良, 王道泽, 汪彦欣, 孙文明, 孔令文, 夏博, 王国迪, 2020. 一种昆虫病原线虫培养瓶. 中国实用新型专利, ZL201921993353.0. 2020-10-09.]
- Chen R, Xu CX, Tao QJ, Zheng Y, Jin LX, Wang DZ, 2019. Effects of 5 insecticides on survivorship and infectivity of entomopathogenic nematodes and their interactions. *China Plant Protection*, 39(12): 14–19. [陈瑞, 徐彩霞, 陶巧静, 郑毅, 金立新, 王道泽, 2019. 5 种杀虫剂对小卷蛾斯氏线虫存活和侵染的影响及其交互作用. 中国植保导刊, 39(12): 14–19.]
- Chen S, Gollop N, Glazer I, 2005. Cross-stress tolerance and expression of stress-related proteins in osmotically desiccated entomopathogenic *Steinernema feltiae* IS-6. *Parasitology*, 131(Pt 5): 695–703.
- Chen SL, Li XH, Yan AH, Spiridonov S, Moens M, 2006a. A new entomopathogenic nematode, *Steinernema hebeiense* sp. n. (Rhabditida: Steinernematidae), from north China. *Nematology*, 8(4): 563–574.
- Chen SB, Glazer I, Gollop N, Cash P, Argo E, Innes A, Stewart E, Davidson I, Wilson MJ, 2006b. Proteomic analysis of the entomopathogenic nematode *Steinernema feltiae* IS-6 IJs under evaporative and osmotic stresses. *Molecular & Biochemical Parasitology*, 145(2): 195–204.
- Chen SB, Yang HW, Jiang SN, 2000. Studies on the biochemical characters of *Steinernema carpocapsae* BJ in anhydrobiosis. *Acta Parasitologica et Medica Entomologica Sinica*, 7(1): 30–34. [陈松笔, 杨怀文, 蒋书楠, 2000. 脱水休眠小卷蛾斯氏线虫的生化特性研究. 寄生虫与医学昆虫学报, 7(1): 30–34.]
- Chen SB, Yang HW, Jiang SN, 2001a. Morphology and oxygen consumption of *Steinernema carpocapsae* in anhydrobiosis. *Acta Entomologica Sinica*, 44(1): 62–66. [陈松笔, 杨怀文, 蒋书楠, 2001a. 小卷蛾线虫脱水休眠的形态与耗氧量. 昆虫学报, 44(1): 62–66.]
- Chen SB, Yang HW, Jiang SN, Jian H, Yang XF, 2001b. Ultrastructure of *Steinernema carpocapsae* BJ in anhydrous dormancy. *Acta Zoologica Sinica*, 47(4): 442–446, 487–488. [陈松笔, 杨怀文, 蒋书楠, 简恒, 杨秀芬, 2001b. 脱水休眠小卷蛾斯氏线虫的超微结构. 动物学报, 47(4): 442–446, 487–488.]
- Chen SB, Yang HW, Luan XQ, Jian H, Yang XF, 2001c. Study on the leaf surface antidesiccator beneficial to entomopathogenic nematode. *Acta Phytophylacica Sinica*, 28(1): 17–22. [陈松笔, 杨怀文, 栾显群, 简恒, 杨秀芬, 2001c. 昆虫病原线虫叶面用抗干燥剂的研究. 植物保护学报, 28(1): 17–22.]
- Chen SB, Yang HW, Luan XQ, Yang XF, Jian H, 1999. The antidesiccation ability of entomopathogenous nematodes. *Chinese Journal of Biological Control*, 15(3): 100–103. [陈松笔, 杨怀文, 栾显群, 杨秀芬, 简恒, 1999. 昆虫病原线虫抗干燥能力研究. 中国生物防治学报, 15(3): 100–103.]
- Chen SL, Li XH, Shi JH, 2006. Occurrence of entomopathogenic nematodes in Hebei Province. *Chinese Journal of Biological Control*, 22(1): 40–44. [陈书龙, 李秀花, 史均环, 2006. 河北省昆虫病原线虫资源调查. 中国生物防治学报, 22(1): 40–44.]
- Chen XQ, Liu QZ, Cao JF, Wei TY, Shi Y, Guo L, 2016. Evaluation of two entomopathogenic nematode species for control of *Batocera lineolata* on walnut trees. *Chinese Journal of Biological Control*, 32(4): 456–461. [陈贤群, 刘奇志, 曹景富, 魏太阳, 施艳, 郭蕾, 2016. 两种昆虫病原线虫防治核桃云斑天牛田间效果评价. 中国生物防治学报, 32(4): 456–461.]
- Chen XR, Xu CX, Han DB, Zhao M, Zhou FC, 2021. Control of *Bemisia tabaci* in pepper by *Steinernema carpocapsae*. *Chinese Journal of Biological Control*, 37(1): 110–116. [陈向荣, 徐彩霞, 韩杜斌, 赵明, 周福才, 2021. 小卷蛾斯氏线虫对辣椒烟粉虱的控制作用. 中国生物防治学报, 37(1): 110–116.]
- Cliche TA, Ensign JC, 2003. For the insect pathogen *Photorhabdus luminescens*, which end of a nematode is out? *Applied Microbiology and Biotechnology*, 69(4): 1890–1897.
- Clausi M, Longo A, Rappazzo G, Tarasco E, Vinciguerra MT, 2011. *Steinernema vulcanicum* n. sp. (Rhabditida: Steinernematidae), a new entomopathogenic nematode species from Sicily (Italy). *Nematology*, 13(4): 409–423.

- Converse V, Grewal PS, 1998. Virulence of entomopathogenic nematodes to the western masked chafer *Cyclocephala hirta* (Coleoptera: Scarabaeidae). *Journal of Economic Entomology*, 91(2): 428–432.
- Cui AN, Jiang X, Zheng YF, Wang H, 2015. Antibiotic activities of the crude extract of entomopathogenic nematodes bacteria to *Fusarium moniliforme*. *Hubei Agricultural Sciences*, 54(4): 862–863, 902. [崔安妮, 姜鑫, 郑毅帆, 王欢, 2015. 昆虫病原线虫共生菌毒素对水稻恶苗病菌的抑菌活性. 湖北农业科学, 54(4): 862–863, 902.]
- Cutler GC, Stock SP, 2003. *Steinernema websteri* sp. n. (Rhabditida: Steinernematidae), a new entomopathogenic nematode from China. *Nematologia Mediterranea*, 31: 215–224.
- Daborn PJ, Waterfield N, Silva CP, Au CPY, Sharma S, French-Constant RH, 2002. A single *Photobacterium* gene, makes caterpillars floppy (mcf), allows *Escherichia coli* to persist within and kill insects. *Proceedings of the National Academy of Sciences of the United States of America*, 99(16): 10742–10747.
- Di B, Qian XJ, Li YY, Liu CZ, 2021. Comparison of two rejuvenation methods on the virulence, fecundity and viability of *Heterorhabditis megidis* and *Steinernema feltiae*. *Chinese Journal of Applied Entomology*, 58(6): 1401–1408. [狄斌, 钱秀娟, 李玉玉, 刘长仲, 2021. 两种复壮方法对昆虫病原线虫致病力、生殖力、存活能力的影响. 应用昆虫学报, 58(6): 1401–1408.]
- Didiza L, Lephot TE, Gray VM, 2021. Morphological and molecular phylogenetic description of *Steinernema batswanae* n. sp. (Rhabditida: Steinernematidae): A new species of an entomopathogenic nematode from south Africa. *Archives of Phytopathology and Plant Protection*, 54(19/20): 1603–1616.
- Ding Q, 2024. Combination usage of entomopathogenic nematodes and botanical spray oil against target insect pest. Master dissertation. Guangzhou: Zhongkai University of Agriculture and Engineering. [丁倩, 2024. 昆虫病原线虫与植物油对害虫的协同防控效果. 硕士学位论文. 广州: 仲恺农业工程学院.]
- Dong YJ, Li XH, Duan JQ, Qin YC, Yang XF, Ren J, Li GY, 2020. Improving the yield of xenocoumacin 1 enabled by in situ product removal. *ACS Omega*, 5(32): 20391–20398.
- Dreyer J, Malan AP, Dicks LMT, 2018. Bacteria of the genus *Xenorhabdus*, a novel source of bioactive compounds. *Frontiers in Microbiology*, 9: 3177.
- Dreyer J, Rautenbach M, Booyens E, van Staden AD, Deane SM, Dicks LT, 2019. *Xenorhabdus khoisaniae* SB10 produces Lys-rich PAX lipopeptides and a xenocoumacin in its antimicrobial complex. *BMC Microbiology*, 19(1): 132.
- Du AQ, Ying TT, Zhou ZY, Yu WC, Hu GG, Luo X, Ma MJ, Yu YL, Wang H, Wei B, 2022. Non-ribosomal peptide biosynthetic potential of the nematode symbiont *Photobacterium*. *Environmental Microbiology Reports*, 14(6): 917–925.
- Duchaud E, Rusniok C, Frangeul L, Buchrieser C, Givaudan A, Taourit S, Bocs S, Boursaux-Eude C, Chandler M, Charles JF, Dassa E, Deroose R, Derzelle S, Freyssinet G, Gaudriault S, Médigue C, Lanois A, Powell K, Signier P, Vincent R, Wingate V, Zouine M, Glaser P, Boemare N, Danchin A, Kunst F, 2003. The genome sequence of the entomopathogenic bacterium *Photobacterium luminescens*. *Nature Biotechnology*, 21(11): 1307–1313.
- Duong DA, Espinosa-Artiles P, Orozco RA, Molnár I, Stock SP, 2019. Draft genome assembly of the entomopathogenic bacterium *Photobacterium luminescens* subsp. *sonorensis* Caborca. *Microbiology Resource Announcements*, 8(36): e00692.
- Eckstein S, Dominelli N, Brachmann A, Heermann R, 2019. Phenotypic heterogeneity of the insect pathogen *Photobacterium luminescens*: Insights into the fate of secondary cells. *Applied and Environmental Microbiology*, 85(22): 1910–1919.
- Ehlers RU, 2001. Mass production of entomopathogenic nematodes for plant protection. *Applied Microbiology and Biotechnology*, 56(5/6): 623–633.
- Ehlers RU, Han RC, 1998. Cultivation of axenic *Heterorhabditis* spp. dauer juveniles and their response to non-specific *Photobacterium luminescens* food signals. *Nematologica*, 44(4): 425–435.
- Ehlers RU, Stoessel S, Wyss U, 1990. The influence of phase variants of *Xenorhabdus* spp. and *Escherichia coli* (Enterobacteriaceae) on the propagation of entomopathogenic nematodes of the genera *Steinernema* and *Heterorhabditis*. *Review of Nematology*, 13: 417–424.
- Fan FF, Zhang ZH, Xu CT, 2015. Pathogenicity of the nematode *S. longicaudum* X-7 against the larvae of *Polyphylla gracilicornis* (a turfgrass underground pest). *Jiangsu Agricultural Sciences*, 43(9): 162–164. [范芳芳, 张宗豪, 徐成体, 2015. *S. longicaudum* X-7 线虫对草坪地下害虫小云斑鳃金龟幼虫的致病力. 江苏农业科学, 43(9): 162–164.]
- Fang FF, Cheng QQ, Li H, Chen XZ, Zhai JH, Hu B, 2019. A powder formulation of entomopathogenic nematodes and its preparation method. Chinese Invention Patent, ZL201710046453.1. 2019–08–13. [方分分, 程清泉, 李欢, 陈新中, 翟金花, 胡彬, 2019. 一种昆虫病原线虫粉剂及其制备方法. 中国发明专利, ZL201710046453.1. 2019–08–13.]
- Fang XL, Zhang WG, Yi XH, Wang YH, Zhang X, 2008. Antimicrobial activity of extracellular metabolites from three symbiotic bacteria of entomopathogenic nematode. *Chinese*

- Journal of Biological Control*, 24(4): 354–358. [方香玲, 张武岗, 易晓华, 王永宏, 张兴, 2008. 三株昆虫病原线虫共生菌胞外产物的抑菌活性. *中国生物防治*, 24(4): 354–358.]
- Feng SP, Han RC, 2005. Advances in storage and formulation of entomopathogenic nematodes *Steinernema* and *Heterorhabditis*. *Plant Protection*, 31(5): 20–25. [冯世鹏, 韩日畴, 2005. 昆虫病原线虫的贮存和剂型研究进展. *植物保护*, 31(5): 20–25.]
- Feng SP, Han RC, Qiu XH, Cao L, Chen JH, Wang GH, 2006. Storage of osmotically treated entomopathogenic nematode *Steinernema carpocapsae*. *Insect Science*, 13(4): 263–269.
- Friedman MJ, 1990. Commercial Production and Development, Entomopathogenic Nematodes in Biological Control. Boca Raton: Chemical Rubber Company Press. 153–172.
- Fu JR, Liu QZ, Li XY, 2018a. Isolation and pathogenicity of symbiotic bacteria associated with the entomopathogenic nematode *Heterorhabditis beicherriana*. *Chinese Journal of Biological Control*, 34(1): 133–140. [付俊瑞, 刘奇志, 李星月, 2018a. 昆虫病原线虫异小杆属新种 *Heterorhabditis beicherriana* 共生菌的分离及致病性. *中国生物防治学报*, 34(1): 133–140.]
- Fu JR, Liu QZ, Peng MX, 2018b. Isolation and pathogenicity of symbiotic bacteria associated with the nematode XJZL1409 (*Heterorhabditis bacteriophora*). *Journal of Environmental Entomology*, 40(5): 1128–1135. [付俊瑞, 刘奇志, 彭梦香, 2018b. 异小杆线虫 XJZL1409 (*Heterorhabditis bacteriophora*) 共生菌的分离及致病性研究. *环境昆虫学报*, 40(5): 1128–1135.]
- Fu ZQ, Li ZX, Xu NF, Qiu SQ, Zhan JH, Du HW, Guo WX, 2020. Preliminary study on the control effect of entomopathogenic nematodes on *Bradysia odoriphaga* in dajingou leek field. *Vegetables*, 2020(9): 64–67. [付在秋, 李朝霞, 许念芳, 邱淑芹, 战金辉, 杜洪伟, 郭文秀, 2020. 应用昆虫病原线虫防治大金钩韭菜田韭蛆效果初探. *蔬菜*, 2020(9): 64–67.]
- Ganguly S, Singh LK, 2000. *Steinernema thermophilum* sp. n. (Rhabditida: Steinernematidae) from India. *International Journal of Nematology*, 10(2): 183–191.
- Gao Y, Zhan F, Chen C, Ba H, Yang R, Wang N, Hou X, Hou M, Shi Y, Long X, 2022. Screening, identification and optimization of fermentation conditions of antagonistic *Xenorhabdus bovienii* 445 against *Aspergillus niger*. *Xinjiang Agricultural Sciences*, 59(5): 1203–1215. [高宇洁, 詹发强, 陈澄, 包慧芳, 杨蓉, 王宁, 侯新强, 侯敏, 史应武, 龙宣杞, 2022. 黑曲霉拮抗菌 *Xenorhabdus bovienii* 445 筛选、鉴定及发酵优化. *新疆农业科学*, 59(5): 1203–1215.]
- Gardner SL, Stock SP, Kaya HK, 1994. A new species of *Heterorhabditis* from the Hawaiian islands. *Journal of Parasitology*, 80(1): 100–106.
- Gaugler R, Brown I, Shapiro-Ilan D, Atwa A, 2002. Automated technology for in vivo mass production of entomopathogenic nematodes. *Biological Control*, 24(2): 199–206.
- Ghazal S, Swanson E, Simpson S, Morris K, Abebe-Akele F, Kelley Thomas W, Khalil KM, Tisa LS, 2017. Permanent draft genome sequence of *Photorhabdus temperata* strain hm, an entomopathogenic bacterium isolated from nematodes. *Genome Announcement*, 5(37): e00974.
- Glaser RW, Fox H, 1930. A nematode parasite of the Japanese beetle (*Popillia japonica* Newm.). *Science*, 71(1827): 16–17.
- Gorgadze O, Lortkipanidze M, Ogier JC, Tailliez P, Burjanadze M, 2015. *Steinernema tbilisiensis* sp. n. (Nematoda: Steinernematidae): A new species of entomopathogenic nematode from Georgia. *Journal of Agricultural Science and Technology*, 5(4): 264–276.
- Grewal PS, Power KT, Grewal SK, Suggars A, Haupricht S, 2004. Enhanced consistency in biological control of white grubs (Coleoptera: Scarabaeidae) with new strains of entomopathogenic nematodes. *Biological Control*, 30(1): 73–82.
- Gu XH, Cui YH, Ruan WB, Cao L, Xue D, Li JZ, Tu YL, Hang DF, Han RC, 2021. Protection efficacy of tobacco seedlings by entomopathogenic nematodes. *Journal of Environmental Entomology*, 43(6): 1529–1535. [谷星慧, 崔永和, 阮维斌, 曹莉, 薛丹, 李江舟, 屠娅玲, 杭德发, 韩日畴, 2021. 昆虫病原线虫对烟苗的保护效果. *环境昆虫学报*, 43(6): 1529–1535.]
- Gu XH, Wang J, Qu L, Li JZ, Cao L, Han RC, Huang ZH, Cui YH, Yang HL, Liu YT, 2023. Use of dimethyl sulfoxide in the preparation of a yield-enhancing formulation for entomopathogenic nematodes. Chinese Invention Patent, ZL202110953231.4. 2023-07-18. [谷星慧, 王杰, 屈玲, 李江舟, 曹莉, 韩日畴, 黄智华, 崔永和, 杨海林, 刘珏廷, 2023. 二甲基亚砜在制备提高昆虫病原线虫产量制剂中的应用. 中国发明专利, ZL202110953231.4. 2023-07-18.]
- Guo WX, Chen H, Zhou XH, Zhai YF, Zheng L, YU Y, Zhao GY, 2019a. A field irrigation-assisted application device for entomopathogenic nematodes. Chinese Utility Model Patent, ZL201821298589.8. 2019-06-07. [郭文秀, 陈浩, 周仙红, 翟一凡, 郑礼, 于毅, 赵国玉, 2019a. 一种昆虫病原线虫田间随水施加装置. 中国实用新型专利, ZL201821298589.8. 2019-06-07.]
- Guo WX, Chen H, Zhou XH, Li LL, Men XY, Yu Y, Zhao GY, 2019b. A field application device for switchable administration of different entomopathogenic nematodes. Chinese Utility Model Patent, ZL201821298264.X. 2019-06-07. [郭文秀, 陈浩, 周仙红, 李丽莉, 门兴元, 于毅, 赵国玉, 2019b. 可切换施用不同

- 昆虫病原线虫的田间施用装置. 中国实用新型专利, ZL201821298264.X. 2019-06-07.]
- Guo WX, Chen H, Zhou XH, Zhang AX, Yu Y, Zhao GY, 2019c. An injection device of entomopathogenic nematodes for controlling longhorn beetles. Chinese Utility Model Patent, ZL201821298975.7. 2019-06-07. [郭文秀, 陈浩, 周仙红, 张安盛, 于毅, 赵国玉, 2019c. 用于防治天牛的昆虫病原线虫注射装置. 中国实用新型专利, ZL201821298975.7. 2019-06-07.]
- Guo WX, Han RC, Yan X, 2020. Use of humus soil as a carrier for storing entomopathogenic nematodes. Chinese Invention Patent, ZL201710297018.6. 2020-10-20. [郭文秀, 韩日畴, 颜珣, 2020. 腐殖土作为载体在贮存昆虫病原线虫中的应用. 中国发明专利, ZL201710297018.6. 2020-10-20.]
- Guo WX, Men XY, Li LL, Lv SH, Yu Y, Cui HY, Song YY, 2024. A moisture-retaining sustained-release capsule for entomopathogenic nematodes in soil and its preparation method. Chinese Invention Patent, ZL202410153781.1. 2024-06-11. [郭文秀, 门兴元, 李丽莉, 吕素洪, 于毅, 崔洪莹, 宋莹莹, 2024. 一种用于土壤中昆虫病原线虫的保湿缓释胶囊及制备方法. 中国发明专利, ZL202410153781.1. 2024-06-11.]
- Guo WX, Zhao GY, Chen H, Zhou XH, Yu Y, 2019d. A culturing chamber for entomopathogenic nematodes. Chinese Utility Model Patent, ZL201821298942.2. 2019-07-16. [郭文秀, 赵国玉, 陈浩, 周仙红, 于毅, 2019d. 一种昆虫病原线虫培养盒. 中国实用新型专利, ZL201821298942.2. 2019-07-16.]
- Guo WX, Zhao GY, Chen H, Zhou XH, Yu Y, 2019e. A packaging box for entomopathogenic nematodes with cultivation function. Chinese Utility Model Patent, ZL201821298622.7. 2019-06-18. [郭文秀, 赵国玉, 周仙红, 陈浩, 于毅, 2019e. 一种具有培养功能的昆虫病原线虫包装盒. 中国实用新型专利, ZL201821298622.7. 2019-06-18.]
- Guo WX, Zhou XH, Chen H, Zhuang QY, Zhang AX, Yu Y, Zhao GY, 2019f. A storage device for maintaining suspension state of entomopathogenic nematodes. Chinese Utility Model Patent, ZL201821298587.9. 2019-06-07. [郭文秀, 周仙红, 陈浩, 庄乾营, 于毅, 赵国玉, 2019f. 一种保持昆虫病原线虫悬浮状态的储存装置. 中国实用新型专利, ZL201821298587.9. 2019-06-07.]
- Guo SQ, Wang ZY, Liu BL, Gao JT, Fang XL, Tang Q, Bilal M, Wang YH, Zhang X, 2019. Effects of *cpxR* on the growth characteristics and antibiotic production of *Xenorhabdus nematophila*. *Microbial Biotechnology*, 12(3): 447–458.
- Guo SQ, Zhang SJ, Fang XL, Liu Q, Gao JT, Bilal M, Wang YH, Zhang X, 2017a. Regulation of antimicrobial activity and xenocoumacins biosynthesis by pH in *Xenorhabdus nematophila*. *Microbial Cell Factories*, 16(1): 203.
- Guo W, Yan X, Zhao G, Han R, 2013. Efficacy of entomopathogenic *Sternernema* and *Heterorhabditis* nematodes against white grubs (Coleoptera: Scarabaeidae) in peanut fields. *Journal of Economic Entomology*, 106(3): 1112–1117.
- Guo WX, Wang XY, Li LL, Lü SH, Cui HY, Song YY, Yu Y, Men XY, 2024. Pathogenicity of five entomopathogenic nematodes to *Tuta absoluta*. *Chinese Journal of Applied Entomology*, 61(4): 741–748. [郭文秀, 王晓钰, 李丽莉, 吕素洪, 崔洪莹, 宋莹莹, 于毅, 门兴元, 2024. 五种昆虫病原线虫对番茄潜叶蛾的致死作用. 应用昆虫学报, 61(4): 741–748.]
- Guo WX, Wang XY, Men XY, Wang CL, Pan HK, Song YY, Cui HY, Lv SH, Yu Y, Li LL, 2023. Field efficacy of *Steinernema carpocapsae* (Rhabditida: Steinernematidae) strain all in the control of fall armyworm, *Spodoptera frugiperda* (Lepidoptera: Noctuidae) in maize. *Biological Control*, 180: 105202.
- Guo WX, Yan X, Han RC, 2017b. Adapted formulations for entomopathogenic nematodes, *Steinernema* and *Heterorhabditis* spp. *Nematology*, 19(5): 587–596.
- Guo WX, Yan X, Zhao GY, Chen JH, Han RC, 2015. Efficacy of entomopathogenic *Steinernema* and *Heterorhabditis* nematodes against *Holotrichia oblita*. *Journal of Pest Science*, 88(2): 359–368.
- Guo WX, Yan X, Zhao GY, Han RC, 2017c. Increased efficacy of entomopathogenic nematode-insecticide combinations against *Holotrichia oblita* (Coleoptera: Scarabaeidae). *Journal of Economic Entomology*, 110(1): 41–51.
- Guo XX, 2020. Screening of resistant entomopathogenic nematodes strains and functional analysis of resistant genes. Master dissertation. Baoding: Hebei Agricultural University. [郭笑笑, 2020. 昆虫病原线虫抗性品系筛选与抗性基因功能分析. 硕士学位论文. 保定: 河北农业大学.]
- Guo XX, Yang Q, Liu SQ, Wang QY, Nangong ZY, 2019. Effect of PirAB toxin from *Xenorhabdus nematophila* on the activities of related enzymes in *Helicoverpa armigera*. *Journal of Agricultural Biotechnology*, 27(7): 1266–1274. [郭笑笑, 杨晴, 刘淑琴, 王勤英, 南宫自艳, 2019. 嗜线虫致病杆菌 PirAB 毒素对棉铃虫相关酶活性的影响. 农业生物技术学报, 27(7): 1266–1274.]
- Ha BJ, Wei XQ, Lu PY, Qing H, Guo JX, Zhang RJ, Chen L, Li XY, Hu B, Wang SY, Xu Y, Fu Z, Shapiro-Ilan D, Ruan WB, 2025. Natural UV protectants and humectants to improve the efficiency of *Steinernema carpocapsae* in controlling foliar pests. *Pest Management Science*, 81(3): 1422–1431.
- Han RC, 1996. The effects of inoculum size on yield of *Steinernema carpocapsae* and *Heterorhabditis bacteriophora* in liquid culture.

- Nematologica*, 42(5): 546–553.
- Han RC, Cao L, He XY, Li QJ, Liu XL, Huang H, Pang Y, He ME, 2000. Recovery response of *Heterorhabditis bacteriophora* and *Steinernema carpocapsae* to different non-symbiotic microorganisms. *Entomologia Sinica*, 7(3): 271–277.
- Han RC, Cao L, Liu XL, 1992. Relationship between medium composition, inoculum size, temperature and culture time in the yields of *Steinernema* and *Heterorhabditis* nematodes. *Fundamental and Applied Nematology*, 15(3): 223–229.
- Han RC, Cao L, Liu XL, 1993. Effects of inoculum size, temperature and time on *in vitro* production of *Steinernema carpocapsae* Agriotes. *Nematologica*, 39(1/4): 366–375.
- Han RC, Ehlers RU, 1999. Trans-specific nematocidal activity of *Photorhabdus luminescens*. *Nematology*, 1(7): 687–693.
- Han RC, Ehlers RU, 2000. Pathogenicity, development and reproduction of *Heterorhabditis bacteriophora* and *Steinernema carpocapsae* under axenic *in vivo* conditions. *Journal of Invertebrate Pathology*, 75(1): 55–58.
- Han RC, Ehlers RU, 2001. Effect of *Photorhabdus luminescens* phase variants on the *in vivo* and *in vitro* development and reproduction of the entomopathogenic nematodes *Heterorhabditis bacteriophora* and *Steinernema carpocapsae*. *FEMS Microbiology Ecology*, 35(3): 239–247.
- Han RC, He XY, Cao L, Liu XL, Yang P, Li QJ, 2000. An attapulgitic clay formulation for storage of entomopathogenic nematode *Steinernema carpocapsae*. *Natural Enemies of Insects*, 22(2): 49–53. [韩日畴, 何向阳, 曹莉, 刘秀玲, 杨平, 李秋剑, 2000. 利用吸附性物质贮存昆虫病原线虫. 昆虫天敌, 22(2): 49–53.]
- Han RC, Leite LG, Yan X, Chacon-Orozco JG, Shapiro-Ilan DI, 2024. Mass production//Shapiro-Ilan DI, Lewis EE (eds.). Entomopathogenic Nematodes as Biological Control Agents. Wallingford: CABI. 182–200.
- Han RC, Li LY, 2003. A liquid culture method for entomopathogenic nematodes. Chinese Invention Patent, ZL99117297.3. 2003-07-23. [韩日畴, 李丽英, 2003. 一种昆虫病原线虫的液体培养方法. 中国发明专利, ZL99117297.3. 2003-07-23.]
- Han RC, Li LY, Pang XF, 1997. Modelling of the culture parameters for production of *Steinernema carpocapsae* and *Heterorhabditis bacteriophora* in solid cultures. *Natural Enemies of Insects*, 19(2): 75–83. [韩日畴, 李丽英, 庞雄飞, 1997. 昆虫病原线虫固体培养系统中主要培养参数的优化. 昆虫天敌, 19(2): 75–83.]
- Han RC, Li LY, Wouts WM, 1990a. The disparities between morphological and functional dimorphism in *Xenorhabdus luminescens*. *Natural Enemies of Insects*, 12(1): 25–32. [韩日畴, 李丽英, Wim M Wouts, 1990a. *Xenorhabdus luminescens* 菌株菌落功能二型性的特殊性. 昆虫天敌, 12(1): 25–32.]
- Han RC, Liu XL, Cao L, 1990b. A modified method of establishing monoxenic culture of entomophagous nematodes *Steinernema* and *Heterorhabditis*. *Natural Enemies of Insects*, 12(3): 113–115. [韩日畴, 刘秀玲, 曹莉, 1990b. 昆虫病原线虫一级种单菌培养的一种方法. 昆虫天敌, 12(3): 113–115.]
- Han RC, Pang XF, Li LY, 1995. Optimization of the medium components for the solid culture of entomopathogenic *Steinernema* and *Heterorhabditis* nematodes. *Natural Enemies of Insects*, 17(4): 153–164. [韩日畴, 庞雄飞, 李丽英, 1995. 昆虫病原线虫固体培养系统干粉培养基的优化. 昆虫天敌, 17(4): 153–164.]
- Han RC, Wouts WM, Li LY, 1990. Development of *Heterorhabditis* spp. strains as characteristics of possible *Xenorhabdus luminescens* subspecies. *Review of Nematology*, 13: 411–415.
- Han YF, Zhang SJ, Wang Y, Gao JT, Han JH, Yan ZQ, Ta YQ, Wang YH, 2024. Enhancing the yield of Xenocoumacin 1 in *Xenorhabdus nematophila* YL001 by optimizing the fermentation process. *Scientific Reports*, 14(1): 13506.
- Han YF, Zhao XT, He MR, Zhang SJ, Tang GJ, Wang YH, 2025. OmpR indirectly regulates biosynthesis of xenocoumacin 1 in *Xenorhabdus nematophila*. *Microorganisms*, 13(6): 1360.
- Han WS, Wang ZR, Liu Y, Gao JL, Zhao DX, Zhong YH, Zhao S, 2021. Effects evaluation of different entomopathogenic nematode strains against *Aethina tumida* (Coleoptera: Nitidulidae). *Chinese Journal of Tropical Crops*, 42(5): 1402–1408. [韩文素, 王泽如, 刘宇, 高景林, 赵冬香, 钟义海, 赵珊, 2021. 不同品系昆虫病原线虫对蜂巢小甲虫的致病力研究. 热带作物学报, 42(5): 1402–1408.]
- Hao J, Yang Q, Li TH, Bai XB, Nangong ZY, Wang QY, 2017. Research on a new solid medium of entomo-pathogenic nematodes and its parameters optimization. *China Plant Protection*, 37(6): 14–19, 65. [郝杰, 杨晴, 李天慧, 白向宾, 南宫自艳, 王勤英, 2017. 一种新的昆虫病原线虫固体培养基及其参数优化研究. 中国植保导刊, 37(6): 14–19, 65.]
- Hartley CJ, Lillis PE, Owens RA, Griffin CT, 2019. Infective juveniles of entomopathogenic nematodes (*Steinernema* and *Heterorhabditis*) secrete ascarosides and respond to interspecific dispersal signals. *Journal of Invertebrate Pathology*, 168: 107257.
- Hatting JL, Moore SD, Malan AP, 2019. Microbial control of phytophagous invertebrate pests in south Africa: Current status and future prospects. *Journal of Invertebrate Pathology*, 165: 54–66.
- Hirao A, Ehlers RU, 2009. Influence of cell density and phase variants of bacterial symbionts (*Xenorhabdus* spp.) on dauer

- juvenile recovery and development of biocontrol nematodes *Steinernema carpocapsae* and *S. feltiae* (Nematoda: Rhabditida). *Applied Microbiology and Biotechnology*, 84(1): 77–85.
- Hsieh FC, Tzeng CY, Tseng JT, Tsai YS, Meng M, Kao SS, 2009. Isolation and characterization of the native entomopathogenic nematode, *Heterorhabditis brevicaudis*, and its symbiotic bacteria from Taiwan. *Current Microbiology*, 58(6): 564–570.
- Hu SB, Liu ZQ, Zhang X, Zhang GY, Xie YL, Ding XZ, Mo XT, Francis Stewart A, Fu J, Zhang YM, Xia LQ, 2016. “Cre/loxP plus BAC”: A strategy for direct cloning of large DNA fragment and its applications in *Photorhabdus luminescens* and *Agrobacterium tumefaciens*. *Scientific Reports*, 6: 29087.
- Hu SL, Wang LT, Qiu XH, Yan X, Zhang JP, Han RC, 2014. Study on the transposon-mutants of *Photorhabdus luminescens* LN2. *Journal of Environmental Entomology*, 36(6): 905–911. [胡素丽, 王立婷, 丘雪红, 颜珣, 张建萍, 韩日畴, 2014. *Photorhabdus luminescens* LN2 转座突变菌株的研究. 环境昆虫学报, 36(6): 905–911.]
- Hua S, Zhao HY, Wang CL, Wang XP, Li CJ, 2018. Infectivity of entomopathogenic nematode to larva of *Lymantria dispar*. *Journal of Northeast Forestry University*, 46(5): 89–92. [滑莎, 赵红盈, 王从丽, 王鑫鹏, 李春杰, 2018. 昆虫病原线虫对舞毒蛾幼虫的侵染能力. 东北林业大学学报, 46(5): 89–92.]
- Huang S, 2023. Determination of insect control effect of entomopathogenic nematode and functional verification of antidesiccation genes. Master dissertation. Zhangjiakou: Hebei North University. [黄山, 2023. 昆虫病原线虫防虫效果测定及抗干燥基因功能验证. 硕士学位论文. 张家口: 河北北方学院.]
- Huang XY, Li C, Zhang K, Li KY, Xie JJ, Peng YY, Quan MF, Sun YJ, Hu YB, Xia LQ, Hu SB, 2024. Function and global regulation of type III secretion system and flagella in entomopathogenic nematode symbiotic bacteria. *International Journal of Molecular Sciences*, 25: 7579.
- Huang YC, Zhu QQ, Zhang X, 2020. Experimental study on biological agents for control of larva of white radish *Phyllotreta striolata*. *Modern Agricultural Science and Technology*, 2020(10): 73, 77. [黄亚川, 朱青青, 张鼎, 2020. 生物药剂防治白萝卜黄曲条跳甲幼虫试验. 现代农业科技, 2020(10): 73, 77.]
- Hunt DJ, Subbotin SA, 2016. Advances in entomopathogenic nematode taxonomy and phylogeny// Hunt DJ, Nguyen KB (eds.). *Nematology Monographs and Perspectives*, Boston LCC. 35.
- Imai Y, Meyer KJ, Iinishi A, Favre-Godal Q, Green R, Manuse S, Caboni M, Mori M, Niles S, Ghiglieri M, Honrao C, Ma XY, Guo JJ, Makriyannis A, Linares-Otaya L, Böhringer N, Wuisan ZG, Kaur H, Wu RR, Mateus A, Typas A, Savitski MM, Espinoza JL, O'Rourke A, Nelson KE, Hiller S, Noinaj N, Schäberle TF, D'Onofrio A, Lewis K, 2019. A new antibiotic selectively kills gram-negative pathogens. *Nature*, 576(7787): 459–464.
- Jeong PY, Jung M, Yim YH, Kim H, Park M, Hong E, Lee W, Kim YH, Kim K, Paik YK, 2005. Chemical structure and biological activity of the *Caenorhabditis elegans* dauer-inducing pheromone. *Nature*, 433(7025): 541–545.
- Jian H, Reid AP, Hunt DJ, 1997. *Steinernema ceratophorum* n. sp. (Nematoda: Steinernematidae) a new entomopathogenic nematode from Northeast China. *Systematic Parasitology*, 37(2): 115–125.
- Jian H, Cao CL, Liu Q, Wang JL, 2011. Insect tissue enzymatic hydrolysate and its application in artificial cultivation of entomopathogenic nematodes. Chinese Invention Patent, ZL200810227286.1. 2011–07–20. [简恒, 曹翠玲, 刘倩, 王金利, 2011. 昆虫组织酶解物及其在昆虫病原线虫人工培养上的应用. 中国发明专利, ZL200810227286.1. 2011–07–20.]
- Jiang F, Shen JW, Cheng JX, Wang X, Yang JG, Li NN, Gao N, Jin Q, 2022. N-terminal signal peptides facilitate the engineering of PVC complex as a potent protein delivery system. *Science Advances*, 8(17): eabm2343.
- Jin LX, Dong DX, Zhang PS, Sun QL, Chen JR, Jiang QF, 2019. Control effect of entomopathogenic nematodes and their combination with insecticides on *Phyllotreta striolata*. *Zhejiang Agricultural Sciences*, 60(3): 443–445. [金立新, 董代率, 张佩胜, 孙泉良, 陈建仁, 蒋群峰, 2019. 昆虫病原线虫及其与药剂混用对青菜黄曲条跳甲的防治效果. 浙江农业科学, 60(3): 443–445.]
- Jin YL, Han RC, 2009. Infectivity of *Steinernema carpocapsae* All strain to the diamondback moth on cabbage leaves. *Plant Protection*, 35(5): 128–131. [金永玲, 韩日畴, 2009. 小卷蛾斯氏线虫 All 品系对小菜蛾感染力测定. 植物保护, 35(5): 128–131.]
- Jin YL, Han RC, Cong B, 2013. Screening of adjuvants for entomopathogenic nematodes of *Steinernema carpocapsae* All strain in foliar application. *Plant Protection*, 39(5): 194–200. [金永玲, 韩日畴, 丛斌, 2013. 昆虫病原线虫 *Steinernema carpocapsae* All 品系叶面应用辅助剂的筛选. 植物保护, 39(5): 194–200.]
- Kaplan F, Alborn HT, von Reuss SH, Ajredini R, Ali JG, Akyazi F, Stelinski LL, Edison AS, Schroeder FC, Teal PE, 2012. Interspecific nematode signals regulate dispersal behavior. *PLoS ONE*, 7(6): e38735.
- Kaya HK, Aguilera MM, Alumai A, Choo HY, de la Torre M, Fodor A, Ganguly S, Hazır S, Lakatos T, Pye A, Wilson M, Yamanaka S, Yang HW, Ehlers RU, 2006. Status of entomopathogenic nematodes and their symbiotic bacteria from selected countries

- or regions of the world. *Biological Control*, 38(1): 134–155.
- Kim K, Sato K, Shibuya M, Zeiger DM, Butcher RA, Ragains JR, Clardy J, Touhara K, Sengupta P, 2009. Two chemoreceptors mediate developmental effects of dauer pheromone in *Caenorhabditis elegans*. *Science*, 326(5955): 994–998.
- Kong XX, Huang ZH, Gu XH, Cui YH, Li JZ, Han RC, Jin YL, Cao L, 2022. Dimethyl sulfoxide and ascarosides improve the growth and yields of entomopathogenic nematodes in liquid cultures. *Journal of Invertebrate Pathology*, 193: 107800.
- Koppenhöfer AM, Brown IM, Gaugler R, Grewal PS, Kaya HK, Klein MG, 2000. Synergism of entomopathogenic nematodes and imidacloprid against white grubs: Greenhouse and field evaluation. *Biological Control*, 19(3): 245–251.
- Koppenhöfer AM, Cowles RS, Cowles EA, Fuzy EM, Baumgartner L, 2002. Comparison of neonicotinoid insecticides as synergists for entomopathogenic nematodes. *Biological Control*, 24(1): 90–97.
- Koppenhöfer AM, Fuzy EM, 2003. *Steinernema scarabaei* for the control of white grubs. *Biological Control*, 28(1): 47–59.
- Koppenhöfer AM, Grewal PS, Fuzy EM, 2006. Virulence of the entomopathogenic nematodes *Heterorhabditis bacteriophora*, *Heterorhabditis zealandica*, and *Steinernema scarabaei* against five white grub species (Coleoptera: Scarabaeidae) of economic importance in turfgrass in North America. *Biological Control*, 38(3): 397–404.
- Kozodoi EM, 1984. New entomopathogenic nematode *Neoaplectana anomali* sp. n. (Rhabditida, Steinernematidae) and observations of its biology. *Zoological Journal*, 63: 1605–1609.
- Labaude S, Griffin CT, 2018. Transmission success of entomopathogenic nematodes used in pest control. *Insects*, 9(2): 72.
- Lanois-Nouri A, Pantel L, Fu J, Houard J, Ogier JC, Polikanov YS, Racine E, Wang HL, Gaudriault S, Givaudan A, Gualtieri M, 2022. The odilorhabdin antibiotic biosynthetic cluster and acetyltransferase self-resistance locus are niche and species specific. *mBio*, 13(1): e0282621.
- Lei LP, Nangong ZY, Li LT, Dong ZP, Wang QY, 2013. Efficacy of entomopathogenic nematodes against *Athetis lepigone*. *Journal of Environmental Entomology*, 35(2): 171–175. [雷利平, 南宫自艳, 李立涛, 董志平, 王勤英, 2013. 昆虫病原线虫对二点委夜蛾致病力的研究. 环境昆虫学报, 35(2): 171–175.]
- Li AH, Zhang Y, Zhang H, Sun F, Zhang XP, Sun RH, 2012. Occurrence dynamics and biocontrol technology of peach and small heartworm in jujube orchard. *Chinese Fruit Trees*, 2012(2): 53–56. [李爱华, 张勇, 张辉, 孙峰, 张学萍, 孙瑞红, 2012. 枣园桃小食心虫的发生动态及生防技术研究. 中国果树, 2012(2): 53–56.]
- Li B, Qiu DW, Wang SC, 2021a. Complete genome sequence data of *Xenorhabdus budapestensis* strain C72, a candidate biological control agent from China. *Plant Disease*, 105(10): 3276–3278.
- Li CJ, Tan GZ, Wang Y, Xu YL, 2011. Occurrence of entomopathogenic nematodes resources and overwintering conditions in Heilongjiang Province. *Plant Protection*, 37(2): 120–123. [李春杰, 谭国忠, 王义, 许艳丽, 2011. 黑龙江省昆虫病原线虫资源和越冬情况调查初报. 植物保护, 37(2): 120–123.]
- Li CJ, Tan GZ, Wang Y, Xu YL, 2014. Optimizing freezing storage conditions for the entomopathogenic nematode isolated from the cold region in China. *Plant Protection*, 40(5): 49–54. [李春杰, 谭国忠, 王义, 许艳丽, 2014. 寒区昆虫病原线虫冷冻储存条件优化. 植物保护, 40(5): 49–54.]
- Li CJ, Wang LF, Pan FJ, Xu YL, 2015. Effect of pretreatment method on freezing survival of entomopathogenic nematodes. *Journal of Northwest A&F University (Natural Science Edition)*, 43(2): 198–202. [李春杰, 王丽芳, 潘凤娟, 许艳丽, 2015. 预处理方式对昆虫病原线虫冷冻存活的影响. 西北农林科技大学学报(自然科学版), 43(2): 198–202.]
- Li CJ, Wang Y, Xu YL, Zhang ZM, Meng J, Yao Q, Song J, 2013. Efficacy of a mixed formulation of *Heterorhabditis bacteriophora* ZT (Heilongjiang isolate) and *Steinernema carpocapsae* All against *Bradysia odoriphaga*. *Chinese Journal of Applied Entomology*, 50(1): 235–241. [李春杰, 王义, 许艳丽, 张志明, 孟洁, 姚钦, 宋洁, 2013. 寒区昆虫病原线虫对韭菜迟眼蕈蚊的防治研究. 应用昆虫学报, 50(1): 235–241.]
- Li CJ, Xu YL, Tan GZ, Zhan LL, He ZT, Lang L, 2008. Laboratory screening of technology for *in vivo* mass production of *Heterorhabditis bacteriophora*. *System Sciences and Comprehensive Studies in Agriculture*, 24(4): 398–402. [李春杰, 许艳丽, 谭国忠, 站丽莉, 何振涛, 郎林, 2008. 异小杆线虫活体繁殖技术筛选. 农业系统科学与综合研究, 24(4): 398–402.]
- Li CJ, Xu YL, Wang Y, Liu CZ, Qian XJ, 2007. Evaluation of the effects of different methods on the quality of entomopathogenic nematodes. *Plant Protection*, 33(5): 88–92. [李春杰, 许艳丽, 王义, 刘长仲, 钱秀娟, 2007. 不同培养方法对昆虫病原线虫质量影响及其评价. 植物保护, 33(5): 88–92.]
- Li CQ, Wang Y, Han RC, Song CG, Ni HP, Liu FT, Li J, Duan XK, Zhang JP, 2011. Determination of pathogenicity of *Steinernema carpocapsae* All to three insect pests. *Journal of Environmental Entomology*, 33(4): 512–516. [李长青, 王媛, 韩日畴, 宋长贵, 倪海平, 刘丰涛, 李姣, 段祥坤, 张建萍, 2011. 昆虫病原线虫 *Steinernema carpocapsae* All 对三种害虫的致病力测定. 环境昆虫学报, 33(4): 512–516.]

- Li ET, Qin JH, Feng HL, Li JQ, Li XF, Nyamwasa I, Cao YZ, Ruan WB, Li KB, Yin J, 2021b. Immune-related genes of the larval *Holotrichia parallela* in response to entomopathogenic nematodes *Heterorhabditis beicheriana* LF. *BMC Genomics*, 22(1): 192.
- Li ET, Cao YZ, Zhang S, Li KB, Li JQ, Li XF, Zhang HR, Yin J, 2019. Control efficacy of the combined application of the entomopathogenic nematode and Bt against the oriental armyworm, *Mythimna separata* (Walker). *Plant Protection*, 45(6): 295–302. [李而涛, 曹雅忠, 张帅, 李克斌, 李金桥, 李晓峰, 张海茹, 尹姣, 2019. 昆虫病原线虫与Bt联用对黏虫的防治效果. 植物保护, 45(6): 295–302.]
- Li ET, Li JY, Li XF, Li JQ, Cao YZ, Zhang S, Li KB, Yin J, 2018. Overview of the enhanced effects of entomopathogenic nematodes to control grubs. The 2018 Annual Academic Conference of the Chinese Society of Plant Protection and the Award Ceremony of the Plant Protection Science and Technology Awards. Xi'an: 199–206. [李而涛, 李建一, 李晓峰, 李金桥, 曹雅忠, 张帅, 李克斌, 尹姣, 2018. 昆虫病原线虫防治蛴螬增效措施概述. 中国植物保护学会 2018 年学术年会暨植保科技奖颁奖大会. 西安: 199–206.]
- Li ET, Li KB, Zhang S, Cao YZ, Lu JJ, Yin J, 2020. Identification and biological characteristics of entomopathogenic nematodes with effective control against white grubs. *Journal of Plant Protection*, 47(5): 1108–1121. [李而涛, 李克斌, 张帅, 曹雅忠, 陆俊姣, 尹姣, 2020. 有效防治蛴螬的昆虫病原线虫鉴定及其生物学特性. 植物保护学报, 47(5): 1108–1121.]
- Li ET, Lu QH, Zhang DF, Kong WJ, An CJ, 2022. Effects of entomopathogenic nematodes on the physiobiochemistry and histopathology of fall armyworm *Spodoptera frugiperda*. *Journal of Plant Protection*, 49(6): 1731–1741. [李而涛, 鲁祺晗, 张丹凤, 孔维捷, 安春菊, 2022. 昆虫病原线虫对草地贪夜蛾生理生化和组织病理学的影响. 植物保护学报, 49(6): 1731–1741.]
- Li ET, Wu HJ, Wang ZM, Li KB, Zhang S, Cao YZ, Yin J, 2023. PI3K/Akt/CncC signaling pathway mediates the response to EPN-Bt infection in *Holotrichia parallela* larvae. *Pest Management Science*, 79(5): 1660–1673.
- Li G, 2017. Biocontrol efficacy of entomopathogenic nematode against larvae of tobacco grubs. Master dissertation. Guangzhou: South China Agricultural University. [李根, 2017. 昆虫病原线虫对烟草蛴螬的控制作用. 硕士学位论文. 广州: 华南农业大学.]
- Li G, Xu WJ, Wang XZ, Hu YX, Xian JD, 2017. Evaluation of different strains entomopathogenic nematodes against *Agrotis ypsilon* Rottemberg (Lepidoptera, Noctuidae) in tobacco. *Journal of Environmental Entomology*, 39(5): 1025–1031. [李根, 许文君, 王新中, 户艳霞, 洗继东, 2017. 不同品系昆虫病原线虫对烟草小地老虎的致病力. 环境昆虫学报, 39(5): 1025–1031.]
- Li JJ, Wei XQ, Pei ZX, Sun J, Xi JL, Li XY, Shapiro-Ilan D, Ruan WB, 2024a. Volatile organic compounds released from entomopathogenic nematode-infected insect cadavers for the biocontrol of *Meloidogyne incognita*. *Pest Management Science*, 80(10): 5400–5411.
- Li JJ, Yu ZQ, Zhang XQ, Ruan WB, 2023. A method for cultivating entomopathogenic nematodes in mealworms and its application. Chinese Invention Patent, ZL202111016372.X. 2023-03-03. [李晶晶, 于真琦, 张秀清, 阮维斌, 2023. 黄粉虫体内培养昆虫病原线虫的方法及其应用. 中国发明专利, ZL202111016372.X. 2023-03-03.]
- Li M, Wu GF, Liu CK, Chen YQ, Qiu LH, Pang Y, 2009. Expression and activity of a probable toxin from *Photorhabdus luminescens*. *Molecular Biology Reports*, 36(4): 785–790.
- Li MG, Ha BJ, Li YC, Vrieling K, Fu Z, Yu QL, Rasmann S, Wei XQ, Ruan WB, 2024b. Toxicological impacts of microplastics on virulence, reproduction and physiological process of entomopathogenic nematodes. *Ecotoxicology and Environmental Safety*, 273: 116153.
- Li N, Wang ZG, Yan AH, Wang QY, Huang DZ, Bi YG, Ma XC, 2011. Anti-desiccation ability of several entomopathogenic nematode strains and their pathogenicity to *Grapholita molesta*. *Journal of Agricultural University of Hebei*, 34(2): 90–93. [李宁, 王志刚, 阎爱华, 王勤英, 黄大庄, 毕拥国, 马向超, 2011. 昆虫病原线虫抗干燥能力及其对梨小食心虫致死能力研究. 河北农业大学学报, 34(2): 90–93.]
- Li XT, Chen H, Wang XD, Qian XJ, 2024. Effects of insulin on the growth and development of the entomopathogenic nematode *Steinernema kraussei* 0657L and its symbiotic bacterium *Xenorhabdus bovienii* 0657L. *Acta Entomologica Sinica*, 67(1): 38–47. [李茜童, 陈红, 王兴铎, 钱秀娟, 2024. 胰岛素对昆虫病原线虫卡森氏线虫 0657L 生长发育及其共生细菌伯氏致病杆菌 0657L 的影响. 昆虫学报, 67(1): 38–47.]
- Li SC, Liang MF, Zhu XM, 1987. Factors affecting the productivity of insect pathogenic nematodes. *Natural Enemies of Insects*, 9(4): 224–227. [李素春, 梁梅芳, 朱雪媚, 1987. 昆虫病原线虫产量与诸因素关系的研究. 昆虫天敌, 9(4): 224–227.]
- Li SH, 2023. Screening of highly pathogenic entomopathogenic nematode strains against meadow moth and study on their application conditions. Master dissertation. Harbin: Northeast Agricultural University. [李朔涵, 2023. 草地螟高致病力昆虫病原线虫品系筛选及施用条件研究. 硕士学位论文. 哈尔滨: 东北农业大学.]

- Li X, Men XY, Wang JH, Lv SH, Li LL, Cui HY, Song YY, Fang XS, Song ZQ, Guo WX, Yu Y, 2023. Curative efficacy of entomopathogenic nematodes against white grubs in honeysuckle fields. *Frontiers in Sustainable Food Systems*, 7: 1155133.
- Li XW, Yi SW, Chen LM, Hafeez M, Zhang ZJ, Zhang JM, Zhou SX, Dong WY, Huang J, Lu YB, 2024c. The application of entomopathogenic nematode modified microbial communities within nesting mounds of the red imported fire ants, *Solenopsis invicta*. *Science of the Total Environment*, 912: 168748.
- Li XF, Wang GH, 1989. Survey on distribution of Steinernematidae and Heterorhabditidae in Fujian, Guangdong and Hainan. *Chinese Journal of Zoology*, 24(3): 1–4, 65. [李小峰, 王国汉, 1989. 闽、粤、海南地区的斯氏线虫科及异小杆线虫科分布调查. *动物学杂志*, 24(3): 1–4, 65.]
- Li XT, Qian XJ, Chen H, Wang XD, Wu X, 2024d. Anti-aging effect of glycerophosphocholine in *Steinernema kraussei* 0657L. *Frontiers in Physiology*, 15: 1346579.
- Li XY, Liu QZ, Nermu J, Půža V, Mráček Z, 2012. *Heterorhabditis beicherriana* n. sp. (Nematoda: Heterorhabditidae), a new entomopathogenic nematode from the Shunyi district of Beijing, China. *Zootaxa*, 3569(1): 25–40.
- Li XL, Zhang FQ, He NY, 2019. Intelligent integrated live culture device for entomopathogenic nematodes. Chinese Utility Model Patent, ZL201821793326.4. 2019-12-03. [李小龙, 张凤琴, 何农跃, 2019. 智能一体化昆虫病原线虫活体培养装置. 中国实用新型专利, ZL201821793326.4. 2019-12-03.]
- Li XL, Zhang FQ, He NY, 2021. A method and culture device for large-scale production of entomopathogenic nematodes. Chinese Invention Patent, ZL201811297686.X. 2021-06-29. [李小龙, 张凤琴, 何农跃, 2021. 一种昆虫病原线虫规模化培养的方法及培养装置. 中国发明专利, ZL201811297686.X. 2021-06-29.]
- Li XY, Zhang HT, Fu HJ, Zeng SC, Yi J, Yang WY, Chen XJ, Chen C, Yi CY, Xu SN, 2025. Preparation method and application of a formulation of entomopathogenic nematodes for controlling sweet potato weevil. Chinese Invention Patent, 202510001064.1. 2025-04-18. [李星月, 张海婷, 符慧娟, 曾钰程, 易军, 杨武云, 陈晓娟, 陈诚, 冀拯宇, 许诗诺, 2025. 一种防治甘薯小象甲的昆虫病原制剂、其制备方法和应用. 中国发明专利, 202510001064.1. 2025-04-18.]
- Li XY, Zhang H, Liu QZ, Yang WY, Li QY, Chen C, Zhu CH, Fu HJ, 2020. An entomopathogenic nematode formulation and its preparation method and application. Chinese Invention Patent, ZL201811491348.X. 2020-09-22. [李星月, 张鸿, 刘奇志, 杨武云, 李其勇, 陈诚, 朱从桦, 符慧娟, 2020. 一种昆虫病原线虫制剂及制备方法和应用. 中国发明专利, ZL201811491348.X. 2020-09-22.]
- Li Y, Zhang GL, Liu Z, Zong J, 1991. Study on vitality of entomopathogenic nematode sprayed with high pressure sprayer. *Chinese Journal of Biological Control*, 7(1): 22–23. [黎彦, 张桂兰, 刘峥, 宗靖, 1991. 用喷雾器喷施线虫对其存活和侵染力的影响. *生物防治通报*, 7(1): 22–23.]
- Liao HJ, Yu HJ, Liao SQ, Li XL, He CX, Hu B, 2023a. A mixing and crushing device for culture media of entomopathogenic nematodes. Chinese Utility Model Patent, ZL202320695378.2. 2023-10-27. [廖慧军, 喻华杰, 廖胜强, 李小龙, 何成新, 胡冰, 2023a. 一种用于昆虫病原线虫培养基混合粉碎的装置. 中国实用新型专利, ZL202320695378.2. 2023-10-27.]
- Liao HJ, Yu HJ, Liao SQ, Li XL, He CX, Hu B, 2023b. A liquid fermentation culture device for entomopathogenic nematodes. Chinese Utility Model Patent, ZL202320439527.9. 2023-08-25. [廖慧军, 喻华杰, 廖胜强, 李小龙, 何成新, 胡冰, 2023b. 一种昆虫病原线虫液体发酵培养装置. 中国实用新型专利, ZL202320439527.9. 2023-08-25.]
- Liao HJ, Yu HJ, Liao SQ, Li XL, He CX, Hu B, 2023c. A storage device for maintaining suspension state of entomopathogenic nematodes. Chinese Utility Model Patent, 202320314970.3. 2023-08-25. [廖慧军, 喻华杰, 廖胜强, 李小龙, 何成新, 胡冰, 2023c. 一种保持昆虫病原线虫悬浮状态的储存装置. 中国实用新型专利, 202320314970.3. 2023-08-25.]
- Liang ML, Li ZY, Dai QX, Lu YY, Cheng KW, Wang L, 2020. Virulence of four entomopathogenic nematodes on fall armyworm, *Spodoptera frugiperda* (J.E. Smith) (Lepidoptera: Noctuidae). *Journal of Biosafety*, 29(2): 82–89. [梁铭荣, 李子园, 戴钊萱, 陆永跃, 陈科伟, 王磊, 2020. 4 种昆虫病原线虫对草地贪夜蛾的致死作用. *生物安全学报*, 29(2): 82–89.]
- Lis M, Sajnaga E, Skowronek M, Wiater A, Rachwał K, Kazimierzczak W, 2021. *Steinernema sandneri* n. sp. (Rhabditida: Steinernematidae), a new entomopathogenic nematode from Poland. *Journal of Nematology*, 53: e2021–51.
- Liu D, Burton S, Glancy T, Li ZS, Hampton R, Meade T, Merlo DJ, 2003. Insect resistance conferred by 283-kDa *Photorhabdus luminescens* protein TcdA in *Arabidopsis thaliana*. *Nature Biotechnology*, 21(10): 1222–1228.
- Liu HL, Robinson DS, Wu ZY, Kuo RT, Yoshikuni Y, Blaby IK, Cheng JF, 2020. Bacterial genome editing by coupling Cre-lox and CRISPR-Cas9 systems. *PLoS ONE*, 15(11): e0241867.
- Liu J, Bai H, Song P, Nangong ZY, Dong ZP, Li ZY, Wang QY, 2022. Insecticidal activity of chitinases from *Xenorhabdus nematophila* HB310 and its relationship with the toxin complex. *Toxins*, 14(9): 646.

- Liu J, NanGong ZY, Zhang J, Song P, Tang Y, Gao Y, Wang QY, 2019. Expression and characterization of two chitinases with synergistic effect and antifungal activity from *Xenorhabdus nematophila*. *World Journal of Microbiology and Biotechnology*, 35(7): 106.
- Liu J, Xue YX, Zang JC, Qian XJ, 2025. Additional investigation of entomopathogenic nematode resources in Gansu Province, northwestern China and screening of the high virulence entomopathogenic nematode strain against *Spodoptera frugiperda* (Lepidoptera: Noctuidae). *Acta Entomologica Sinica*, 68(4): 432–449. [刘佳, 薛银霞, 臧建成, 钱秀娟, 2025. 甘肃省昆虫病原线虫资源补充调查及对草地贪夜蛾高致病力线虫品系筛选. *昆虫学报*, 68(4): 432–449.]
- Liu JQ, Li WW, Wang CL, Liang H, Chu ZN, Wang GP, Jin LH, Guo WX, Yu Y, 2025. Mixtures of entomopathogenic nematodes provided successful suppression of *Bactrocera tau* by stimulated dispersal, aggregation, and invasion. *Biological Control*, 208: 105858.
- Liu M, Lin Y, Zhang JM, 2025. Pathogenicity evaluation of four pathogenic nematodes on *Spodoptera exigua* at pupal stage. *Journal of Zhejiang Agricultural Sciences*, 66(4): 1001–1004. [刘敏, 林雅, 章金明, 2025. 4 种病原线虫对蛹期甜菜夜蛾的致病力评价. *浙江农业科学*, 66(4): 1001–1004.]
- Liu NX, Liang MF, Zhang ZY, Zheng LS, Liang YQ, 1990. Research on application techniques of *Steinernema glaseri* 85011 for controlling bamboo weevils and its compatibility with pesticides. *Natural Enemies of Insects*, 12(3): 142–148. [刘南欣, 梁梅芳, 张振英, 郑莲生, 梁永奇, 1990. 格氏线虫中国品系 (85011) 防治竹象虫的施用技术及与农药混用的研究. *昆虫天敌*, 12(3): 142–148.]
- Liu Q, Han GJ, Li CM, Huang LX, Lu YR, Xia Y, Zhang N, Xu J, 2024. Control effect of entomopathogenic nematode N-Yz1 to rice stem borer, *Chilo suppressalis*. *Chinese Journal of Biological Control*, 40(5): 1015–1021. [刘琴, 韩光杰, 李传明, 黄立鑫, 陆玉荣, 夏杨, 张楠, 徐健, 2024. 昆虫病原线虫 N-Yz1 对水稻二化螟的控制作用. *中国生物防治学报*, 40(5): 1015–1021.]
- Liu Q, Xu J, Li CM, Han GJ, Huang LX, Lu YR, Xia Y, 2023. A culture method of *Steinernema* spp., pupae capsule formulation and its preparation method. Chinese Invention Patent, ZL202210642552.7. 2023-10-13. [刘琴, 徐健, 李传明, 韩光杰, 黄立鑫, 陆玉荣, 夏杨, 2023. 斯氏线虫的培养方法、斯氏线虫蛹胶囊制剂及其制备方法. 中国发明专利, ZL202210642552.7. 2023-10-13.]
- Liu QZ, Zhang LJ, Wang YZ, Du XK, Sun HY, Mráček Z, 2009. Bio-analysis on EPNs of *Steinernema longicaudum* BPS strain. *Chinese Agricultural Science Bulletin*, 25(2): 236–239. [刘奇志, 张丽娟, 王玉柱, 杜小康, 孙浩元, Zdeněk Mráček, 2009. 昆虫病原线虫 (*Steinernema longicaudum* BPS 品系) 生物学测定. *中国农学通报*, 25(2): 236–239.]
- Liu WT, Chen TL, Hou RF, Chen CC, Tu WC, 2020. The invasion and encapsulation of the entomopathogenic nematode, *Steinernema abbasi*, in *Aedes albopictus* (Diptera: Culicidae) larvae. *Insects*, 11(12): 832.
- Liu X, Li Q, Wang YH, Xu X, Zhang X, 2006. The antifungal activity of secondary metabolic products from *Xenorhabdus nematophilus* YL001. *Journal of Plant Protection*, 33(3): 277–281. [刘霞, 李骞, 王永宏, 许贤, 张兴, 2006. 嗜线虫致病杆菌 YL001 菌株代谢产物的抑菌活性. *植物保护学报*, 33(3): 277–281.]
- Liu XS, Zhao CM, Wei GS, Wu XR, Yan JF, 1993. Application of entomopathogenic nematodes for controlling soil-emerging larvae of *Carposina sasakii*. *Journal of Hebei Normal University of Science & Technology*, 7(3): 77–78. [刘新生, 赵春明, 魏国树, 吴学仁, 严景芬, 1993. 应用昆虫病原线虫防治桃小食心虫出土期幼虫的研究. *河北农业技术师范学院学报*, 7(3): 77–78.]
- Liu XX, Rao ZC, Han RC, Jin LH, Cao L, 2024. Genes involved in DMSO-mediated yield increase of entomopathogenic nematodes. *Scientific Reports*, 14(1): 31670.
- Liu Y, 2020. Cultivation and application of entomopathogenic nematodes and study on the effects of abiotic factors on it. Master dissertation. Zhuzhou: Hunan University of Technology. [刘勇, 2020. 昆虫病原线虫的培养应用及非生物因子对其影响的研究. 硕士学位论文. 株洲: 湖南工业大学.]
- Liu Y, Zhang YZ, Xu J, Lu X, Miao XX, Duan DY, Xiao Y, Pan Y, 2025. Screening of dominant strains of entomopathogenic nematodes and their control effect on *Chilo suppressalis*. *Hybrid Rice*, 40(1): 85–89. [刘洋, 张玉烛, 许隽, 鲁湘, 苗雪雪, 段德勇, 肖阳, 盘毅, 2025. 昆虫病原线虫优势品系筛选及其对水稻二化螟的防效研究. *杂交水稻*, 40(1): 85–89.]
- Liu Z, Li Y, Zong J, Zhang GL, Chai F, 1998. Studies on the application of entomopathogenic nematode CZ-88 for control of peach fruit-moth. *Journal of Fruit Science*, 15(1): 53–57. [刘峥, 黎彦, 宗靖, 张桂兰, 柴枫, 1998. 利用昆虫病原线虫 CZ-88 防治桃小食心虫的研究. *果树科学*, 15(1): 53–57.]
- Liu J, 1994. A new species of *Heterorhabditis* (Rhabditida: Heterorhabditidae) from China. *Acta Zoologica Sinica*, 19(3): 268–272. [刘杰, 1994. 中国异小杆线虫属一新种(小杆目: 异小杆线虫科). *动物分类学报*, 19(3): 268–272.]
- Lu CK, Xu ZC, Jia FY, Lu YQ, 2004. Laboratory virulence and

- infectivity of *Steinernema carpocapsae* to *Holcocerus hippophaecolus*. *Chinese Journal of Biological Control*, 20(4): 280–282. [路常宽, 许志春, 贾峰勇, 骆有庆, 2004. 小卷蛾斯氏线虫对沙棘木蠹蛾幼虫的室内侵染能力. *中国生物防治学报*, 20(4): 280–282.]
- Lunau S, Stoessel S, Schmidt-Peisker AJ, Ehlers RU, 1993. Establishment of monoxenic inocula for scaling up in vitro cultures of the entomopathogenic nematodes *Steinernema* spp. and *Heterorhabditis* spp.. *Nematologica*, 39(1/4): 385–399.
- Lv CJ, Meng TG, Zhong BZ, Shang ZQ, Li CX, Zahra AA, Abdelrahman TM, 2025. Pathogenicity of *Steinernema carpocapsae* All entomopathogenic nematodes and their symbiotic bacteria as a biological control agent on red palm weevil. *Microorganisms*, 13(5): 971.
- Ma J, Chen SL, Zou Y, Li XH, Han RC, de Clercq P, Moens M, 2010. Natural occurrence of entomopathogenic nematodes in north China. *Russian Journal of Nematology*, 18: 117–126.
- Ma J, Chen SL, De Clercq P, Han RC, Moens M, 2012a. *Steinernema changbaiense* sp. n. (Rhabditida: Steinernematidae), a new species of entomopathogenic nematode from northeast China. *Russian Journal of Nematology*, 20: 97–112.
- Ma J, Chen SL, De Clercq P, Waeyenberge L, Han RC, Moens M, 2012b. A new entomopathogenic nematode, *Steinernema xinbinense* n. sp. (Nematoda: Steinernematidae), from north China. *Nematology*, 14(6): 723–739.
- Ma J, Chen SL, Li XH, Han RC, Khatri-Chhetri HB, De Clercq P, Moens M, 2012c. A new entomopathogenic nematode, *Steinernema tielingense* n. sp. (Rhabditida: Steinernematidae), from north China. *Nematology*, 14(3): 321–338.
- Ma J, Chen SL, Moens M, De Clercq P, Li XH, Han RC, 2013a. Characterization in biological traits of entomopathogenic nematodes isolated from north China. *Journal of Invertebrate Pathology*, 114(3): 268–276.
- Ma J, Chen SL, Moens M, Han RC, De Clercq P, 2013b. Efficacy of entomopathogenic nematodes (Rhabditida: Steinernematidae and Heterorhabditidae) against the chive gnat, *Bradysia odoriphaga*. *Journal of Pest Science*, 86(3): 551–561.
- Ma J, Guo XX, Li XH, Wang RY, Gao B, Chen SL, 2022. Tolerance of different entomopathogenic nematodes to environmental stresses. *Chinese Journal of Biological Control*, 38(1): 188–195. [马娟, 郭笑笑, 李秀花, 王容燕, 高波, 陈书龙, 2022. 不同品系昆虫病原线虫抗性分析. *中国生物防治学报*, 38(1): 188–195.]
- Ma XY, 2016. Study on the preservation materials of entomopathogenic nematodes. Master dissertation. Wuhan: Central China Normal University. [马小雅, 2016. 昆虫病原线虫保存材料的研究. 硕士学位论文. 武汉: 华中师范大学.]
- Machado RAR, Muller A, Ghazal SM, Thanwisai A, Pagès S, Bode HB, Hussein MA, Khalil KM, Tisa LS, 2021. *Photorhabdus heterorhabditis* subsp. *aluminescens* subsp. nov., *Photorhabdus heterorhabditis* subsp. *heterorhabditis* subsp. nov., *Photorhabdus australis* subsp. *thailandensis* subsp. nov., *Photorhabdus australis* subsp. *australis* subsp. nov., and *Photorhabdus aegyptia* sp. nov. isolated from *Heterorhabditis* entomopathogenic nematodes. *International Journal of Systematic and Evolutionary Microbiology*, 71(1): 4610.
- Machado RAR, Abolafia J, Robles MC, Ruiz-Cuenca AN, Bhat AH, Shokoohi E, Půža V, Zhang X, Erb M, Robert CAM, Hibbard B, 2025. Description of *Heterorhabditis americana* n. sp. (Rhabditida, Heterorhabditidae), a new entomopathogenic nematode species isolated in north America. *Parasites and Vectors*, 18(1): 101.
- Machado RAR, Bhat AH, Abolafia J, Shokoohi E, Fallet P, Turlings TCJ, Tarasco E, Půža V, Kajuga J, Yan X, Toepfer S, 2022. *Steinernema africanum* n. sp. (Rhabditida, Steinernematidae), a new entomopathogenic nematode species isolated in the republic of Rwanda. *Journal of Nematology*, 54(1): 20220049.
- Mahmood S, Kumar M, Kumari P, Mahapatro GK, Banerjee N, Sarin NB, 2020. Novel insecticidal chitinase from the insect pathogen *Xenorhabdus nematophila*. *International Journal of Biological Macromolecules*, 159: 394–401.
- Mathur C, Phani V, Kushwah J, Somvanshi VS, Dutta TK, 2019. TcaB, an insecticidal protein from *Photorhabdus akhurstii* causes cytotoxicity in the greater wax moth, *Galleria mellonella*. *Pesticide Biochemistry and Physiology*, 157: 219–229.
- Meesil W, Bode HB, Rückert-Reed C, Shi YM, Pidot SJ, Muangpat P, Rattanarojpong T, Chantratita N, Sitthisak S, Vitta A, Thanwisai A, 2025. Genomic analysis for the identification of bioactive compounds in *Xenorhabdus stockiae* strain RT25.5. *Scientific Reports*, 15(1): 23672.
- Mollah MMI, Dekebo A, Kim Y, 2020. Immunosuppressive activities of novel PLA₂ inhibitors from *Xenorhabdus hominickii*, an entomopathogenic bacterium. *Insects*, 11(8): 505.
- Mráček Z, Liu QZ, Nguyen KB, 2009. *Steinernema xueshanense* n. sp. (Rhabditida, Steinernematidae), a new species of entomopathogenic nematode from the province of Yunnan, southeast Tibetan Mts., China. *Journal of Invertebrate Pathology*, 102(1): 69–78.
- Mráček Z, Nguyen KB, Tailliez P, Boemare N, Chen SL, 2006. *Steinernema sichuanense* n. sp. (Rhabditida, Steinernematidae), a new species of entomopathogenic nematode from the province of Sichuan, east Tibetan Mts., China. *Journal of Invertebrate*

- Pathology*, 93(3): 157–169.
- Muangpat P, Suwannaroj M, Yimthin T, Fukruksa C, Sitthisak S, Chantratita N, Vitta A, Thanwisai A, 2020. Antibacterial activity of *Xenorhabdus* and *Photorhabdus* isolated from entomopathogenic nematodes against antibiotic-resistant bacteria. *PLoS ONE*, 15(6): e0234129.
- Nangong ZY, Guo XX, Yang Q, Song P, Wang QY, Parajulee MN, 2020. Identification of arylphorin interacting with the insecticidal protein PirAB from *Xenorhabdus nematophila* by yeast two-hybrid system. *World Journal of Microbiology and Biotechnology*, 36(4): 56.
- Nangong ZY, Li TH, Yuan J, Li M, Zhang AB, Ren LX, Zhang LN, Zang HM, Wang XJ, Wang QY, 2022. A powder formulation containing entomopathogenic nematodes. Chinese Invention Patent, ZL201910267874.6. 2022-02-11. [南宮自艷, 李天慧, 袁景, 李敏, 张安邦, 任莉鑫, 张丽娜, 臧昊明, 王晓洁, 王勤英, 2022. 一种含有昆虫病原线虫的粉剂. 中国发明专利, ZL201910267874.6. 2022-02-11.]
- Neubacher N, Tobias NJ, Huber M, Cai XF, Glatter T, Pidot SJ, Stinear TP, Lütticke AL, Papenfort K, Bode HB, 2020. Symbiosis, virulence and natural-product biosynthesis in entomopathogenic bacteria are regulated by a small RNA. *Nature Microbiology*, 5(12): 1481–1489.
- Nguyen KB, Pūza V, Mráček Z, 2008a. *Steinernema cholashanense* n. sp. (Rhabditida, Steinernematidae) a new species of entomopathogenic nematode from the province of Sichuan, Chola Shan Mountains, China. *Journal of Invertebrate Pathology*, 97(3): 251–264.
- Nguyen KB, Qiu LH, Zhou Y, Pang Y, 2006. *Steinernema leizhouense* sp. n. (Nematoda: Steinernematidae), a new entomopathogenic nematode from Southern China. *Russian Journal of Nematology*, 14: 101–118.
- Nguyen KB, Shapiro-Ilan DI, Mbata G, 2008b. *Heterorhabditis georgiana* n. sp. (Rhabditida: Heterorhabditidae) from Georgia, USA. *Nematology*, 10(3): 433–448.
- Nie ZY, Yu Q, Song HL, Liu X, Zhang LJ, Geng Y, Yu J, 2023. Effect of high pathogenicity entomopathogenic nematode on nutrient and protective enzymes in *Dioryctria abietella*. *Journal of Northeast Forestry University*, 51(10): 74–79. [聂振业, 于琪, 宋昊霖, 刘行, 张丽君, 耿瑶, 宇佳, 2023. 高致病性病原线虫对冷杉梢斑蛾营养物及体内活性酶的影响. 东北林业大学学报, 51(10): 74–79.]
- Oliveira-Hofman C, Kaplan F, Stevens G, Lewis E, Wu SH, Alborn HT, Perret-Gentil A, Shapiro-Ilan DI, 2019. Pheromone extracts act as boosters for entomopathogenic nematodes efficacy. *Journal of Invertebrate Pathology*, 164: 38–42.
- Pan FJ, Zhang SJ, Chen Y, Meng J, Wu DB, Wang J, Xiao L, Xu YL, 2013. Management of entomopathogenic nematode on *Bradysia odoriphaga* in Ningxia. *Anhui Agricultural Science Bulletin*, 19(3): 83–86. [潘凤娟, 张思佳, 陈延, 蒙静, 武东波, 王佳, 肖亮, 许艳丽, 2013. 利用昆虫病原线虫防治迟眼蕈蚊的研究. 安徽农学通报, 19(3): 83–86.]
- Pan HY, Ding L, Wang YC, Yang J, 1995. Improve of culture medium for entomopathogenic nematodes. *Journal of Jilin Agricultural University*, 17(4): 16–19, 23. [潘洪玉, 丁利, 王迎春, 杨军, 1995. 昆虫病原线虫培养基的筛选和改进. 吉林农业大学学报, 17(4): 16–19, 23.]
- Pan YH, Yu BZ, Berg OG, Jain MK, Bahnson BJ, 2002. Crystal structure of phospholipase A2 complex with the hydrolysis products of platelet activating factor: Equilibrium binding of fatty acid and lysophospholipid-ether at the active site may be mutually exclusive. *Biochemistry*, 41(50): 14790–14800.
- Patil J, Linga V, Mhatre PH, Gowda MT, Rangasamy VS, Pūza V, 2023. *Steinernema indicum* n. sp., a new entomopathogenic nematode (Nematoda: Steinernematidae) from India. *Nematology*, 25(7): 815–833.
- Peters A, 2013. Application and commercialization of nematodes. *Applied Microbiology Biotechnology*, 97(14): 6181–6188.
- Peters A, Han RC, Yan X, Leite LG, 2017. Production of entomopathogenic nematodes//Lacey LA (ed.). *Microbial Control of Insect and Mite Pests: From Theory to Practice*. Amsterdam: Elsevier Academic Press. 157–170.
- Plichta KL, Joyce SA, Clarke D, Waterfield N, Stock SP, 2009. *Heterorhabditis gerrardi* n. sp. (Nematoda: Heterorhabditidae): The hidden host of *Photorhabdus asymbiotica* (Enterobacteriaceae: γ -Proteobacteria). *Journal of Helminthology*, 83(4): 309–320.
- Pūza V, Mráček Z, 2005. Seasonal dynamics of entomopathogenic nematodes of the genera *Steinernema* and *Heterorhabditis* as a response to abiotic factors and abundance of insect hosts. *Journal of Invertebrate Pathology*, 89(2): 116–122.
- Qian XJ, Sun D, Zang JC, Lewis EE, 2024. Effects of insulin on growth, fecundity and pathogenicity of entomopathogenic nematodes. *Chinese Journal of Biological Control*, 40(4): 856–865. [钱秀娟, 孙丹, 臧建成, Edwin E Lewis, 2024. 胰岛素对昆虫病原线虫生长发育、生殖力及致病力的影响. 中国生物防治学报, 40(4): 856–865.]
- Qian XJ, Gu LN, Xing YF, Liu CZ, 2014. Occurrence of entomopathogenic nematodes in Gansu Province. *Acta Agrestia Sinica*, 22(3): 593–599. [钱秀娟, 谷黎娜, 邢玉芳, 刘长仲, 2014. 甘肃省昆虫病原线虫区系研究. 草地学报, 22(3): 593–599.]

- Qian XJ, Xu YL, Wang Y, Liu CZ, Li CJ, Zhao FP, 2005. Study on the virulence of entomogenous nematodes to the larvae of soybean underground pests of north eastern beetles. *Soybean Science*, 24(3): 224–228. [钱秀娟, 许艳丽, Wang Yi, 刘长仲, 李春杰, 赵风平, 2005. 昆虫病原线虫对大豆地下害虫东北大黑鳃金龟幼虫的致病力研究. *大豆科学*, 24(3): 224–228.]
- Qian XJ, Zhang SP, Xie P, Liu CZ, 2019a. Effects of UV-B radiation on entomopathogenic nematode survival and pathogenicity. *Chinese Journal of Eco-Agriculture*, 27(6): 836–844. [钱秀娟, 张世鹏, 谢攀, 刘长仲, 2019a. 紫外辐射对 9 个昆虫病原线虫品系存活率和致病力的影响. *中国生态农业学报(中英文)*, 27(6): 836–844.]
- Qian XJ, Zhang WL, Liu CZ, 2019b. Virulence of entomopathogenic nematodes and imidacloprid to *Bradysia odoriphaga*. *Journal of Gansu Agricultural University*, 54(2): 102–108. [钱秀娟, 张文丽, 刘长仲, 2019b. 昆虫病原线虫与吡虫啉混用对异迟眼蕈蚊幼虫的毒力测定. *甘肃农业大学学报*, 54(2): 102–108.]
- Qiu LH, Fang YY, Zhou Y, Pang Y, Nguyen KB, 2004. *Steinernema guangdongense* sp. n. (Nematoda: Steinernematidae), a new entomopathogenic nematode from southern China with a note on *S. serratum* (nomen nudum). *Zootaxa*, 704: 1–20.
- Qiu LH, Hu X, Zhou Y, Mei S, Nguyen KB, Pang Y, 2005a. *Steinernema akhursti* sp. n. (Nematoda: Steinernematidae) from Yunnan, China. *Journal of Invertebrate Pathology*, 90(3): 151–160.
- Qiu LH, Hu XQ, Zhou Y, Pang Y, Nguyen KB, 2005b. *Steinernema beddingi* n. sp. (Nematoda: Steinernematidae), a new entomopathogenic nematode from Yunnan, China. *Nematology*, 7(5): 737–749.
- Qiu LH, Yan Y, Zhou Y, Nguyen KB, Pang Y, 2005c. *Steinernema aciari* sp. n. (Nematoda: Steinernematidae), a new entomopathogenic nematode from Guangdong, China. *Journal of Invertebrate Pathology*, 88(1): 58–69.
- Qiu LH, Zhao JX, Wu ZD, Lv ZY, Pang Y, 2011. *Steinernema pui* sp. n. (Rhabditida, Steinernematidae), a new entomopathogenic nematode from Yunnan, China. *Zootaxa*, 2767(1): 1–13.
- Qiu LH, Bedding RA, 2002. Characteristics of protectant synthesis of infective juveniles of *Steinernema carpocapsae* and importance of glycerol as a protectant for survival of the nematodes during osmotic dehydration. *Comparative Biochemistry and Physiology Part B*, 131(4): 757–765.
- Qiu LH, Lacey MJ, Bedding RA, 2000. Permeability of the infective juveniles of *Steinernema carpocapsae* to glycerol during osmotic dehydration and its effect on biochemical adaptation and energy metabolism. *Comparative Biochemistry and Physiology Part B*, 125(3): 411–419.
- Qiu XH, Han RC, Xu ZF, 2004. Biochemical properties of artificial nematode-symbiotic bacterium combinations. *Natural Enemies of Insects*, 26(2): 49–59. [丘雪红, 韩日畴, 许再福, 2004. 不同昆虫病原线虫——共生细菌组合的生化特性研究. *昆虫天敌*, 26(2): 49–59.]
- Qiu XH, Han RC, Yan X, Liu MX, Cao L, Yoshiga T, Kondo E, 2009. Identification and characterization of a novel gene involved in the trans-specific nematocidal activity of *Photorhabdus luminescens* LN2. *Applied and Environmental Microbiology*, 75(12): 4221–4223.
- Qiu XH, Wu CY, Cao L, Ehlers RU, Han RC, 2016. *Photorhabdus luminescens* LN2 requires rpoS for nematocidal activity and nematode development. *FEMS Microbiology Letters*, 363(6): fnw035.
- Qiu XH, Zhan ZB, Yan X, Han RC, 2014. Draft genome sequence and annotation of the entomopathogenic bacterium *Photorhabdus luminescens* LN2, which shows nematocidal activity against *Heterorhabditis bacteriophora* H06 nematodes. *Genome Announcements*, 2(6): e01268–14.
- Ruan WB, 2023. A separation device for entomopathogenic nematodes. Chinese Utility Model Patent, ZL202222834476.8. 2023-04-04. [阮维斌, 2023. 一种昆虫病原线虫分离装置. 中国实用新型专利, ZL202222834476.8. 2023-04-04.]
- Sanda NB, Hou YM, 2023. The symbiotic bacteria-*Xenorhabdus nematophila* All and *Photorhabdus luminescens* H06 strongly affected the phenoloxidase activation of nipa palm hispid, *Octodonta nipae* (Coleoptera: Chrysomelidae) larvae. *Pathogens*, 12(4): 506.
- Sanda NB, Muhammad A, Ali H, Hou YM, 2018. Entomopathogenic nematode *Steinernema carpocapsae* surpasses the cellular immune responses of the hispid beetle, *Octodonta nipae* (Coleoptera: Chrysomelidae). *Microbial Pathogenesis*, 124: 337–345.
- Santhi VS, Ment D, Faigenboim A, Salame L, Soroker V, Hetzroni A, Glazer I, 2021. Behavioral and molecular response of the insect parasitic nematode *Steinernema carpocapsae* to cues emitted by a host, the red palm weevil, *Rhynchophorus ferrugineus*. *Molecular and Biochemical Parasitology*, 241: 111345.
- Shahina F, Tabassum KA, Salma J, Mehreen G, Knoetze R, 2017. *Heterorhabditis pakistanense* n. sp. (Nematoda: Heterorhabditidae) a new entomopathogenic nematode from Pakistan. *Journal of Helminthology*, 91(2): 222–235.
- Shankhu PY, Mathur C, Mandal A, Sagar D, Somvanshi VS, Dutta TK, 2020. Txp40, a protein from *Photorhabdus akhurstii*, conferred potent insecticidal activity against the larvae of

- Helicoverpa armigera*, *Spodoptera litura* and *S. exigua*. *Pest Management Science*, 76(6): 2004–2014.
- Shapiro-Ilan DI, Hazir S, Glazer I, 2017. Microbial Agents for Control of Insect Pests: From Discovery to Commercial Development and Use. Pittsburgh: Academic Press. 91–105.
- Shen CP, Wang GH, 1992. Description and Studies of An Entomopathogenic Nematode: *Steinernema longicaudum* sp. nov.. Beijing: China Science and Technology Press. 220–231. [沈长朋, 王国汉, 1992. 昆虫病原线虫新种 *Steinernema longicaudum* sp. nov. 的描述与研究. 北京: 中国科技出版社. 220–231.]
- Shi YM, Bode HB, 2018. Chemical language and warfare of bacterial natural products in bacteria-nematode-insect interactions. *Natural Product Reports*, 35(4): 309–335.
- Shi YM, Brachmann AO, Westphalen MA, Neubacher N, Tobias NJ, Bode HB, 2019. Dual phenazine gene clusters enable diversification during biosynthesis. *Nature Chemical Biology*, 15(4): 331–339.
- Sicard M, Ferdy JB, Pagès S, Le Brun N, Godelle B, Boemare N, Mouliat C, 2004. When mutualists are pathogens: An experimental study of the symbioses between *Steinernema* (entomopathogenic nematodes) and *Xenorhabdus* (bacteria). *Journal of Evolutionary Biology*, 17(5): 985–993.
- Somvanshi VS, Dubay B, Kushwah J, Ramamoorthy S, Vishnu US, Sankarasubramanian J, Rajendhran J, Rao U, 2019. Draft genome sequences for five *Photorhabdus* bacterial symbionts of entomopathogenic *Heterorhabditis* nematodes isolated from India. *Microbiology Resource Announcement*, 8(4): 1404–1418.
- Song N, Chen LH, Ren XM, Waterfield NR, Yang J, Yang GW, 2021. N-Glycans and sulfated glycosaminoglycans contribute to the action of diverse Tc toxins on mammalian cells. *PLoS Pathogens*, 17(2): e1009244.
- Soni S, Patil J, Linga V, Mhatre PH, Gowda MT, Ganguli J, Pūza V, 2023. *Steinernema shori* n. sp., a new entomopathogenic nematode (Nematoda: Steinernematidae) from India. *Journal of Helminthology*, 97: e72.
- Stock SP, 1996. Pathogenicity of the entomopathogenic nematode *Heterorhabditis argentinensis* (Nematoda: Heterorhabditidae) on insects associated with alfalfa crops. *Revista de la Sociedad Entomológica Argentina*, 55(1/4): 175–177.
- Stock SP, 2019. Partners in crime: Symbiont-assisted resource acquisition in *Steinernema* entomopathogenic nematodes. *Current Opinion in Insect Science*, 32: 22–27.
- Stock SP, Rivera-Orduño B, Flores-Lara Y, 2009. *Heterorhabditis sonorensis* n. sp. (Nematoda: Heterorhabditidae), a natural pathogen of the seasonal cicada *Diceroprocta ornea* (Walker) (Homoptera: Cicadidae) in the Sonoran desert. *Journal of Invertebrate Pathology*, 100(3): 175–184.
- Stock SP, Strong D, Gardner SL, 1996. Identification of *Heterorhabditis* (Nematoda: Heterorhabditidae) from California with a new species isolated from the larvae of the ghost moth *Hepialis californicus* (Lepidoptera: Hepialidae) from the Bodega Bay Natural Reserve. *Fundamental and applied Nematology*, 19(6): 585–592.
- Strauch O, Ehlers RU, 1998. Food signal production of *Photorhabdus luminescens* inducing the recovery of entomopathogenic nematodes *Heterorhabditis* spp. in liquid culture. *Applied Microbiology and Biotechnology*, 50(3): 369–374.
- Sun BJ, Zhang XQ, Song L, Zheng LX, Wei XQ, Gu XH, Cui YH, Hu B, Yoshiga T, Abd-Elgawad MM, Ruan WB, 2021. Evaluation of indigenous entomopathogenic nematodes in southwest China as potential biocontrol agents against *Spodoptera litura* (Lepidoptera: Noctuidae). *Journal of Nematology*, 53: e2021–83.
- Sun LN, Zhang HJ, Yan WT, Yue Q, Li YY, Qiu GS, 2018. Research progress on *Carposina sasakii*. *China Fruits*, 2018(1): 76–81. [孙丽娜, 张怀江, 闫文涛, 岳强, 李艳艳, 仇贵生, 2018. 桃小食心虫研究进展. 中国果树, 2018(1): 76–81.]
- Sun RH, Li AH, 2011. A washer for entomopathogenic nematodes. Chinese Utility Model Patent, ZL201020248472.6. 2011-03-16. [孙瑞红, 李爱华, 2011. 一种昆虫病原线虫清洗器. 中国实用新型专利, ZL201020248472.6. 2011-03-16.]
- Sun RH, Li AH, Han RC, Cao L, Liu XL, 2004. Factors affecting the control of *Bradysina odoriphaga* with entomopathogenic nematode *Heterorhabditis indica* LN2. *Journal of Environmental Entomology*, 26(4): 150–155. [孙瑞红, 李爱华, 韩日畴, 曹莉, 刘秀玲, 2004. 昆虫病原线虫 *Heterorhabditis indica* LN2 品系防治韭菜迟眼蕈蚊的影响因素研究. 昆虫天敌, 26(4): 150–155.]
- Sun YW, Zhang GY, Hou XQ, Xiao S, Yang X, Xie YL, Huang XY, Wang F, Mo XT, Ding XZ, Xia LQ, Hu SB, 2019. SrfABC toxin from *Xenorhabdus stockiae* induces cytotoxicity and apoptosis in HeLa cells. *Toxins*, 11(12): 685.
- Tang P, Wang ZZ, Wu Q, Liu YQ, Shi M, Huang JH, Chen XX, 2019. Natural enemy resources and its application in biological control of *Spodoptera frugiperda*. *Chinese Journal of Applied Entomology*, 56(3): 370–381. [唐璞, 王知知, 吴琼, 刘银泉, 时敏, 黄健华, 陈学新, 2019. 草地贪夜蛾的天敌资源及其生物防治中的应用. 应用昆虫学报, 56(3): 370–381.]
- Tian CL, Zhang JH, Zhu F, Han RC, Li MH, 2024. Research and application progresses of artificial culture technologies for entomopathogenic nematodes. *Acta Entomologica Sinica*, 67(6): 872–884. [田成丽, 张金花, 朱峰, 韩日畴, 李茂海, 2024. 昆

- 虫病原线虫人工培养技术研究与应用进展. 昆虫学报, 67(6): 872–884.]
- Tian CL, Zhu F, Li XY, Zhang JH, Půža V, Shapiro-Ilan D, Zhao D, Liu JW, Zhou JJ, Ding Y, Wang JC, Ma J, Zhu XF, Li MH, Li JP, 2022. *Steinernema populi* n. sp. (Panagrolaimomorpha, Steinernematidae), a new entomopathogenic nematode species from China. *Journal of Helminthology*, 96: e57.
- Tian HP, Liang HZ, Li XY, Chen Q, Ning SH, Hou ZR, 2013. Mass production of entomopathogenic nematodes. *Forest Pest and Disease*, 32(2): 12–13. [田会鹏, 梁洪柱, 李学燕, 陈倩, 宁少华, 侯峥嵘, 2013. 昆虫病原线虫大量培养技术研究. 中国森林病虫, 32(2): 12–13.]
- Tobias NJ, Shi YM, Bode HB, 2018. Refining the natural product repertoire in entomopathogenic bacteria. *Trends in Microbiology*, 26(10): 833–840.
- Tobias NJ, Wolff H, Djahanschiri B, Grundmann F, Kronenwerth M, Shi YM, Simonyi S, Grün P, Shapiro-Ilan S, Pidot SJ, Stinear TP, Ebersberger I, Bode HB, 2017. Natural product diversity associated with the nematode symbionts *Photorhabdus* and *Xenorhabdus*. *Nature Microbiology*, 2(12): 1676–1685.
- Tseng CT, Hou RF, Tang LC, 2018. *Steinernema taiwanensis* n. sp. (Rhabditida: Steinernematidae), a new entomopathogenic nematode from Taiwan. *Zootaxa*, 4434(3): 466–480.
- Visschedyk D, Rochon A, Tempel W, Dimov S, Park HW, Merrill AR, 2012. Certhra toxin, an anthrax-related ADP-ribosyltransferase from *Bacillus cereus*. *Journal of Biological Chemistry*, 287(49): 41089–41102.
- Wang B, Meng FY, Song SQ, Xie B, Jia SX, Xiu DY, Li XP, 2025. Multi-omics analysis of *Curculio dieckmanni* (Coleoptera: Curculionidae) larvae reveals host responses to *Steinernema carpocapsae* infection. *Insects*, 16(5): 503.
- Wang GH, Han RC, Chen JH, Cao L, 2007. Combined efficacy of entomopathogenic nematode *Steinernema carpocapsae* All and pesticide against *Rhabdoscelus lineaticollis* (Heller). *Chinese Journal of Biological Control*, 23(3): 218–222. [王果红, 韩日畴, 陈镜华, 曹莉, 2007. 利用昆虫病原线虫与化学农药混用防治褐纹甘蔗象. 中国生物防治学报, 23(3): 218–222.]
- Wang H, Cong B, Dong H, Qian HT, 2007. Evaluation of insecticidal activities of crude toxins extracted from symbiotic bacteria to *Plutella xylostella* and *Laphygma exigua*. *Chinese Vegetables*, 2007(3): 24–26. [王欢, 丛斌, 董辉, 钱海涛, 2007. 共生菌杀虫毒素对小菜蛾和甜菜夜蛾的杀虫活性. 中国蔬菜, 2007(3): 24–26.]
- Wang H, Cong B, Dong H, Liu YQ, Yang J, 2009. Evaluation of the insecticidal and antibiotic activities of the crude toxins extracted from *Xenorhabdus* and *Photorhabdus* to *Ostrinia furnacalis* and *Bipolaris maydis*. *Journal of Maize Sciences*, 17(6): 109–111. [王欢, 丛斌, 董辉, 刘彦群, 杨吉, 2009. 共生菌毒素对亚洲玉米螟和玉米小斑菌杀虫抑菌活性的测定. 玉米科学, 17(6): 109–111.]
- Wang H, Luan JB, Dong H, Qian HT, Cong B, 2014. Natural occurrence of entomopathogenic nematodes in Liaoning (northeast China). *Journal of Asia-Pacific Entomology*, 17(3): 399–406.
- Wang H, Zeng YQ, Li TJ, Zong XG, Li L, 1998. Study on indoor artificial propagation of entomopathogenic nematodes. *Shandong Journal of Animal Science and Veterinary Medicine*, 19(2): 4–5. [王慧, 曾勇庆, 李铁坚, 宗希阁, 栗磊, 1998. 昆虫病原线虫的室内人工繁殖研究. 山东畜牧兽医, 19(2): 4–5.]
- Wang HC, Lin SJ, Wang HC, Kumar R, Le PT, Leu JH, 2023. A bacterial binary toxin system that kills both insects and aquatic crustaceans: *Photorhabdus* insect-related toxins A and B. *PLoS Pathogens*, 19(5): e1011330.
- Wang J, Cao L, Huang ZH, Gu XH, Cui YH, Li JZ, Li YL, Xu CT, Han RC, 2022. Influence of the ascarosides on the recovery, yield and dispersal of entomopathogenic nematodes. *Journal of Invertebrate Pathology*, 188: 107717.
- Wang J, Dai K, Kong XX, Cao L, Qu L, Jin YL, Li YL, Gu XH, Li JZ, Xu CT, Han RC, 2021. Research progress and perspective on entomopathogenic nematodes. *Journal of Environmental Entomology*, 43(4): 811–839. [王杰, 戴康, 孔祥鑫, 曹莉, 屈玲, 金永玲, 李玉玲, 谷星慧, 李江舟, 徐成体, 韩日畴, 2021. 昆虫病原线虫与环境生物、非生物因素关系的研究进展. 环境昆虫学报, 43(4): 811–839.]
- Wang LF, Li CJ, Xu YL, 2010. Technology for *in vitro* mass production of *Heterorhabditis bacteriophora*-1. *System Sciences and Comprehensive Studies in Agriculture*, 26(1): 31–35. [王丽芳, 李春杰, 许艳丽, 2010. 异小杆线虫离体繁殖技术研究. 农业系统科学与综合研究, 26(1): 31–35.]
- Wang QL, Cui JY, Li ZX, Liu CQ, 1995. Effect of soil moisture on the motion of *Steinernema glaseri* (Nematoda: Steinernematidae). *Chinese Journal of Biological Control*, 11(4): 7–9. [王庆雷, 崔景岳, 李仲秀, 刘春芹, 1995. 不同含水量的土壤对格氏线虫扩散的影响. 中国生物防治学报, 11(4): 7–9.]
- Wang SY, 2022. Investigation of entomopathogenic nematode resources and screening of its fine strains in northern China. Master dissertation. Beijing: Chinese Academy of Agricultural Sciences. [王思雨, 2022. 华北北部地区昆虫病原线虫资源调查及优良品系筛选. 硕士学位论文. 北京: 中国农业科学院.]
- Wang X, Cheng JX, Shen JW, Liu LG, Li NN, Gao N, Jiang F, Jin Q, 2022. Characterization of *Photorhabdus* virulence cassette as a

- causative agent in the emerging pathogen *Photorhabdus asymbiotica*. *Science China Life Sciences*, 65(3): 618–630.
- Wang XP, 2018. The effect and mechanism of biological bacteria against root knot nematodes. Master dissertation. Beijing: Chinese Academy of Sciences. [王鑫鹏, 2018. 根结线虫生防细菌的作用效果及机制初探. 硕士学位论文. 北京: 中国科学院大学.]
- Wang Y, Zhang FY, Wang C, Guo PF, Han YQ, Zhang YT, Sun BJ, Shan SJ, Ruan WB, Pan J, 2022. Antifungal substances produced by *Xenorhabdus bovienii* and its inhibition mechanism against *Fusarium solani*. *International Journal of Molecular Sciences*, 23(16): 9040.
- Wang YD, Xiao C, Yin J, Cao YZ, Li KB, 2012. Evaluation of the impacts on infection ability of entomopathogenic nematodes to grub, *Holotrichia parallela* for three chemical pesticides. *Chinese Journal of Biological Control*, 28(1): 67–73. [王玉东, 肖春, 尹姣, 曹雅忠, 李克斌, 2012. 三种化学杀虫剂对病原线虫侵染暗黑鳃金龟能力的影响. 中国生物防治学报, 28(1): 67–73.]
- Wang YH, Zhang X, 2007. Studies on antimicrobial activity of metabolites of symbiotic bacteria associated with entomopathogenic nematodes. *Journal of Northwest A&F University*, 35(2): 125–130. [王永宏, 张兴, 2007. 2 株昆虫病原线虫共生菌代谢产物的抑菌作用. 西北农林科技大学学报, 35(2): 125–130.]
- Wang YY, Zhang XT, Feng X, Wang X, Jin Q, Jiang F, 2024. Purification of *Photorhabdus* virulence cassette (PVC) protein complexes from *Escherichia coli* for artificial translocation of heterologous cargo proteins. *Bio-protocol*, 14(7): e4966.
- Wang Z, Zhou XJ, Zhu LY, He XC, Jiang L, 2019. Control effect of entomopathogenic nematodes against *Curculio chinensis* Chevrolat larvae. *Journal of Zhejiang Agricultural Sciences*, 60(10): 1768–1769. [王轶, 周小军, 朱丽燕, 何晓婵, 江丽, 2019. 昆虫病原线虫对茶籽象甲幼虫的防治效果. 浙江农业科学, 60(10): 1768–1769.]
- Wang ZW, Liao YM, Zhang GY, Yang CZ, Li ZM, 1999. Culturation of the entomopathogenic nematode and determination of its pathogenicity to two species of the longicorn. *Journal of Guangxi Agricultural and Biological Science*, 18(2): 109–113. [王忠文, 廖咏梅, 张桂英, 杨崇振, 李钟明, 1999. 昆虫病原线虫的培养及其对两种天牛的致病性测定. 广西农业生物科学, 18(2): 109–113.]
- Waterfield N, Kamita SG, Hammock BD, French-Constant R, 2005. The *Photorhabdus* Pir toxins are similar to a developmentally regulated insect protein but show no juvenile hormone esterase activity. *FEMS Microbiology Letters*, 245(1): 47–52.
- Wen X, Nangong ZY, Zhang Y, Song P, Cao KQ, Wang QY, 2012. Antifungal activity of metabolites from *Xenorhabdus nematophila* and *Photorhabdus luminescens* against apple pathogenic fungi. *China Plant Protection*, 32(1): 13–16. [文旭, 南宫自艳, 张园, 宋萍, 曹克强, 王勤英, 2012. 2 种昆虫病原线虫共生菌的代谢产物对苹果病原菌真菌的抑菌作用. 中国植保导刊, 32(1): 13–16.]
- Wesche F, Adihou H, Kaiser A, Wurglics M, Schubert-Zsilavecz M, Kaiser M, Bode HB, 2018. Combined approach of backbone amide linking and on-resin N-methylation for the synthesis of bioactive and metabolically stable peptides. *Journal of Medicinal Chemistry*, 61(9): 3930–3938.
- Wu GQ, Yi YH, Lv YY, Li M, Wang J, Qiu LH, 2015a. The lipopolysaccharide (LPS) of *Photorhabdus luminescens* TT01 can elicit dose and time-dependent immune priming in *Galleria mellonella* larvae. *Journal of Invertebrate Pathology*, 127: 63–72.
- Wu GQ, Zhao ZY, Liu CL, Qiu LH, 2014. Priming *Galleria mellonella* (Lepidoptera: Pyralidae) larvae with heat-killed bacterial cells induced an enhanced immune protection against *Photorhabdus luminescens* TT01 and the role of innate immunity in the process. *Journal of Economic Entomology*, 107(2): 559–569.
- Wu GQ, Yi YH, 2016. Haemocoel injection of PirA₁B₁ to *Galleria mellonella* larvae leads to disruption of the haemocyte immune functions. *Scientific Reports*, 6: 34996.
- Wu HB, Fan K, Xin L, Cao GP, Sun RH, 2015b. Virulence and control effect of entomopathogenic nematodes against *Agrotis ypsilon* (Rotttemberg). *Journal of Plant Protection*, 42(2): 244–250. [武海斌, 范昆, 辛力, 曹广平, 孙瑞红, 2015b. 昆虫病原线虫对小地老虎的致病力测定及防治效果. 植物保护学报, 42(2): 244–250.]
- Wu HB, Gong QT, Chen ZZ, Jiang LL, Gong Y, Xu YY, Sun RH, 2018. Bioefficacy of the combined application of entomopathogenic nematodes and thiamethoxam and its effects on the protective and detoxification enzyme activities in *Bradysia odoriphaga* (Diptera: Sciaridae) larvae. *Entomologica Sinica*, 61(7): 851–859. [武海斌, 宫庆涛, 陈珍珍, 姜莉莉, 公义, 许永玉, 孙瑞红, 2018. 昆虫病原线虫和噻虫嗪混用对韭蛆的杀虫效果及其体内保护酶和解毒酶活性的影响. 昆虫学报, 61(7): 851–859.]
- Wu HB, Liu HF, Gong QT, Zhang KP, Sun RH, 2015c. Application of entomopathogenic nematodes against subterranean pests and influencing factors. *Deciduous Fruits*, 47(5): 24–27. [武海斌, 刘海峰, 宫庆涛, 张坤鹏, 孙瑞红, 2015c. 昆虫病原线虫防治地下害虫的应用及影响因子. 落叶果树, 47(5): 24–27.]
- Wu LH, Wang YT, Hsieh FC, Hsieh C, 2020. Insecticidal activity of *Photorhabdus luminescens* 0805-P2R against *Plutella xylostella*.

- Applied Biochemistry and Biotechnology*, 191(1): 191–200.
- Wu LL, Xie YF, Chang DD, Huang MH, Jiang Y, Qin RF, Jiang D, Wei LL, Wang X, Zhao YN, Dou YW, Wang CL, Li CJ, 2024. Control effect of symbiotic bacteria and their metabolites from entomopathogenic nematode on *Meloidogyne* spp.. *Soils and Crops*, 13(3): 381–389. [吴黎黎, 谢倚帆, 常豆豆, 黄铭慧, 姜野, 秦瑞峰, 蒋丹, 韦柳利, 王旋, 赵亚男, 窦悦文, 王从丽, 李春杰, 2024. 昆虫病原线虫共生菌及代谢产物对根结线虫的防治作用. *土壤与作物*, 13(3): 381–389.]
- Wu SH, Mechrez G, Ment D, Toews MD, Ananth Mani K, Amar Feldbaum R, Shapiro-Ilan DI, 2023a. Tolerance of *Steinernema carpocapsae* infective juveniles in novel nanoparticle formulations to ultraviolet radiation. *Journal of Invertebrate Pathology*, 196: 107851.
- Wu SY, El-Borai FE, Graham JH, Duncan LW, 2018. The saprophytic fungus *Fusarium solani* increases the insecticidal efficacy of the entomopathogenic nematode *Steinernema diaprepesi*. *Journal of Invertebrate Pathology*, 159: 87–94.
- Wu SY, Tang HT, Ban SM, Wen RQ, Mao XF, Deng HH, Abbas MK, Salem SM, Hou YM, 2024. Integrated approaches for *Solenopsis invicta* (Hymenoptera: Formicidae) management: Insights from laboratory studies with entomopathogenic nematodes and insecticides. *Pest Management Science*, 80(9): 4410–4416.
- Wu SY, Tang HT, Zhang C, Tang FX, Lin JH, Wang YX, Chen LH, Hou YM, 2023b. Potential of entomopathogenic nematode-infected insect cadavers for the biocontrol of the red imported fire ant *Solenopsis invicta*. *Pest Management Science*, 79(11): 4383–4389.
- Wu WD, Sun HY, Xi JH, Yin J, Zhang S, Cao YZ, Li KB, Xiao C, 2015d. Observations of the ultrastructure of the fat body and midgut tissues of two white grub species, *Holotrichia parallela* and *H. oblita* (Coleoptera: Melolonthidae), infected by entomopathogenic nematode *Heterorhabditis bacteriophora*. *Acta Entomologica Sinica*, 58(8): 836–845. [吴文丹, 孙昊雨, 席景会, 尹姣, 张帅, 曹雅忠, 李克斌, 肖春, 2015d. 嗜菌异小杆线虫感染后暗黑鳃金龟和大黑鳃金龟幼虫脂肪体和中肠组织超微结构观察. *昆虫学报*, 58(8): 836–845.]
- Wu WD, Yin J, Cao YZ, Xiao C, Li KB, 2014. Entomopathogenic nematodes and their application in China. *Chinese Journal of Biological Control*, 30(6): 817–822. [吴文丹, 尹姣, 曹雅忠, 肖春, 李克斌, 2014. 我国昆虫病原线虫的研究与应用现状. *中国生物防治学报*, 30(6): 817–822.]
- Wu X, 2023. Response of energy substances of *Steinernema kraussei* 0657L and *Heterorhabditis brevicaudis* 0641TY to low humidity stress and the key genes in metabolism. Master dissertation. Lanzhou: Gansu Agricultural University. [吴霞, 2023. 2 种昆虫病原线虫能量物质对低湿胁迫的响应及代谢关键基因研究. 硕士学位论文. 兰州: 甘肃农业大学.]
- Wu X, Li XL, Qian XJ, 2021. Infection of *Galleria mellonella* by entomopathogenic nematodes *Heterorhabditis megidis* 0627M and *Steinernema feltiae* 0619HT. *Chinese Journal of Applied Entomology*, 58(6): 1417–1425. [吴霞, 李小玲, 钱秀娟, 2021. 昆虫病原线虫感染大蜡螟的行为观测. *应用昆虫学报*, 58(6): 1417–1425.]
- Wu X, Zhu XF, Ma CL, Yang YX, Lü XF, Ma HP, Liu CZ, Qian XJ, 2019. Effects of anhydrobiosis on entomopathogenic nematode and its biological characteristics. *Grassland and Turf*, 39(1): 90–94. [吴霞, 朱小芳, 马彩玲, 杨亚贤, 吕小芳, 马宏萍, 刘长仲, 钱秀娟, 2019. 低湿胁迫对昆虫病原线虫生物学特性的影响. *草原与草坪*, 39(1): 90–94.]
- Xi XD, Lu XZ, Zhang XD, Bi YH, Li XC, Yu ZG, 2019. Two novel cyclic depsipeptides Xenematides F and G from the entomopathogenic bacterium *Xenorhabdus budapestensis*. *The Journal of Antibiotics*, 72(10): 736–743.
- Xiao JF, Han LL, Zhao KJ, Hao ZR, Zhu L, Shi ZH, Gao LT, Chen J, 2021. Effects of different media on the resistance of *Steinernema litorale* to ultraviolet radiation. *Plant Protection*, 47(3): 115–121. [肖建飞, 韩岚岚, 赵奎军, 郝子茹, 朱琳, 师正浩, 高丽瞳, 陈娟, 2021. 不同介质对海滨斯氏线虫抗紫外胁迫的影响. *植物保护*, 47(3): 115–121.]
- Xiao YH, 2018. Evaluation of biocontrol potential and optimization of sponge storage conditions of entomopathogenic nematode *Steinernema costaricense*. Master dissertation. Nanjing: Nanjing Agricultural University. [肖宇辉, 2018. 昆虫病原线虫 *Steinernema costaricense* 的生防潜力评价与其海绵存储条件优化. 硕士学位论文. 南京: 南京农业大学.]
- Xie YF, Wang XD, Zhong WH, Zhu DH, He Z, 2020. Transcriptome profile changes associated with heat shock reaction in the entomopathogenic nematode, *Steinernema carpocapsae*. *Frontiers in Physiology*, 11: 721.
- Xing YF, Qian XJ, Liu CZ, Li HG, 2009. Anti-desiccation ability of several entomopathogenic nematodes in Gansu Province. *Guizhou Agricultural Sciences*, 37(1): 90–92. [邢玉芳, 钱秀娟, 刘长仲, 李宏刚, 2009. 甘肃省几种昆虫病原线虫抗干燥能力测定. *贵州农业科学*, 37(1): 90–92.]
- Xu CQ, Xu J, Chen J, Yu J, Chen RM, Qin RM, Xu R, Zhou F, Cheng HZ, 2010. Lethal effects of entomopathogenic nematodes on larvae of *Dorystenes hydropicus* in laboratory experiment. *Chinese Journal of Traditional Chinese Medicine*, 35(10): 1239–1241. [徐常青, 徐江, 陈君, 于晶, 陈荣敏, 覃荣敏, 徐

- 荣, 周峰, 程惠珍, 2010. 昆虫病原线虫对曲牙土天牛室内致死效应初步研究. 中国中药杂志, 35(10): 1239–1241.]
- Xu JL, 1989. Entomopathogenic nematodes *Steinernema* and their symbiotic bacteria. *Entomological Knowledge*, 26(2): 121–123, 99. [徐洁莲, 1989. 昆虫病原斯氏线虫与其联系的共生细菌. 昆虫知识, 26(2): 121–123, 99.]
- Xu JL, Liu XL, 1997. The effects of temperatures, storage methods and moisture carrier on the survival and infectivity of entomopathogenic nematodes. *Natural Enemies of Insects*, 19(3): 97–102. [徐洁莲, 刘秀玲, 1997. 温度、储备方法和载体湿度对斯氏线虫存活和感染力的影响. 昆虫天敌, 19(3): 97–102.]
- Xu JL, Xie RC, 1992. Biological comparisons for two species of entomopathogenic nematodes in wax moth and artificial medium. *Natural Enemies of Insects*, 14(2): 76–82. [徐洁莲, 谢汝创, 1992. 以大蜡螟与人工培养基连续传代的两种斯氏线虫生物学特性比较. 昆虫天敌, 14(2): 76–82.]
- Xu JL, Yang P, 2000. Evaluation of increasing effect using entomopathogenic nematode with binder. *Natural Enemies of Insects*, 22(1): 6–10. [徐洁莲, 杨平, 2000. 昆虫病原线虫与粘合剂混用的增效试验. 昆虫天敌, 22(1): 6–10.]
- Xu YL, Wang Y, Li CJ, Pan FJ, 2018. A sterilizing and mixing device for solid culture of entomopathogenic nematodes. Chinese Invention Patent, ZL201510883160.X. 2018-02-09. [许艳丽, 王义, 李春杰, 潘凤娟, 2018. 一种昆虫病原线虫固体繁殖的灭菌搅拌装置. 中国发明专利, ZL201510883160.X. 2018-02-09.]
- Xu ZF, Wang HG, Li XF, 1991. A new species of the genus *Steinernema* (Rhabditida: Steinernematidae). *Zoological Research*, 12(1): 17–20.
- Yadav M, Rathore JS, 2020. The *hipBA^{Xn}* operon from *Xenorhabdus nematophila* functions as a bonafide toxin-antitoxin module. *Applied Microbiology and Biotechnology*, 104(7): 3081–3095.
- Yan AH, Wang ZG, Jia JC, Bi YG, 2015. A device for large-scale in vivo culture of entomopathogenic nematodes in laboratory. Chinese Utility Model Patent, ZL201420762615.3. 2015-07-15. [阎爱华, 王志刚, 贾金才, 毕拥国, 2015. 一种用于实验室活体大量繁殖昆虫病原线虫的装置. 中国实用新型专利, ZL201420762615.3. 2015-07-15.]
- Yan K, Luo ZL, Hu FL, Tian HS, Yan JC, 2017. Field experiment on controlling the oriental fruit moth with entomopathogenic nematodes. *Journal of Agricultural Research and Application*, 30(3): 58–59, 67. [严凯, 罗泽丽, 胡芳丽, 田华松, 严金成, 2017. 昆虫病原线虫防治梨小食心虫田间试验. 农业研究与应用, 30(3): 58–59, 67.]
- Yan X, Chen GM, Chen YQ, Sun BJ, Gu XH, Ruan WB, Han RC, 2021. Virulence of *Steinernema ceratophorum* against different pest insects and their potential for *in vivo* and *in vitro* culture. *Journal of Nematology*, 53: e2021–46.
- Yan X, De Clercq P, Han RC, Jones J, Chen SL, Moens M, 2011. Osmotic responses of different strains of *Steinernema carpocapsae*. *Nematology*, 13(7): 845–851.
- Yan X, Han RC, 2010. Anhydrobiosis of entomopathogenic nematodes. *Chinese Journal of Biological Control*, 26(1): 90–95. [颜珣, 韩日畴, 2010. 昆虫病原线虫的脱水休眠. 中国生物防治学报, 26(1): 90–95.]
- Yan X, Han RC, 2016. Production technology of the bio-insecticides: Entomopathogenic nematodes. *Journal of Environmental Entomology*, 38(5): 1044–1051. [颜珣, 韩日畴, 2016. 生物杀虫剂-昆虫病原线虫的培养技术. 环境昆虫学报, 38(5): 1044–1051.]
- Yan X, Han RC, Chen SL, Moens M, De Clercq P, 2012. Differential gene expression after osmotic induction in various species of entomopathogenic nematodes. *Russian Journal of Nematology*, 20(2): 167–170.
- Yan X, Han RC, Moens M, Chen SL, De Clercq P, 2013. Field evaluation of entomopathogenic nematodes for biological control of striped flea beetle, *Phyllotreta striolata* (Coleoptera: Chrysomelidae). *BioControl*, 58(2): 247–256.
- Yan X, Lin YN, Huang ZM, Han RC, 2018. Characterisation of biological and biocontrol traits of entomopathogenic nematodes promising for control of striped flea beetle (*Phyllotreta striolata*). *Nematology*, 20(6): 503–518.
- Yan X, Liu P, Chen YQ, 2025. In vitro solid mass production of entomopathogenic nematodes. International Congress on Invertebrate Pathology and Microbial Control & 57th Annual Meeting of the Society for Invertebrate Pathology. Puerto Varas: 63.
- Yan X, Liu XJ, Han RC, Chen SL, De Clercq P, Moens M, 2010. Osmotic induction of anhydrobiosis in entomopathogenic nematodes of the genera *Heterorhabditis* and *Steinernema*. *Biological Control*, 53(3): 325–330.
- Yan X, Shahid Arain M, Lin YY, Gu XH, Zhang LM, Li JZ, Han RC, 2020. Efficacy of entomopathogenic nematodes against the tobacco cutworm, *Spodoptera litura* (Lepidoptera: Noctuidae). *Journal of Economic Entomology*, 113(1): 64–72.
- Yan X, Wang X, Han RC, Qiu XH, 2014. Utilisation of entomopathogenic nematodes, *Heterorhabditis* spp. and *Steinernema* spp. for the control of *Agrotis ipsilon* (Lepidoptera, Noctuidae) in China. *Nematology*, 16(1): 31–40.
- Yan X, Zhao GY, Han RC, 2019. Integrated management of chive gnats (*Bradysia odoriphaga* Yang & Zhang) in chives using entomopathogenic nematodes and low-toxicity insecticides.

- Insects*, 10(6): 161.
- Yang BJ, Yang HW, Liu Z, Yang XF, Yuan JJ, 2007. Comparison of toxicity between extracellular and intracellular proteins of entomopathogenic nematode symbiotic bacteria against *Helicoverpa armigera*. *Chinese Journal of Biological Control*, 23(1): 33–38. [杨保军, 杨怀文, 刘峥, 杨秀芬, 袁京京, 2007. 昆虫病原线虫共生菌胞内外蛋白的提取及杀虫活性比较. 中国生物防治学报, 23(1): 33–38.]
- Yang D, Jang WD, Lee SY, 2021. Production of carminic acid by metabolically engineered *Escherichia coli*. *Journal of the American Chemical Society*, 143(14): 5364–5377.
- Yang FL, Cao TT, Zhang B, Wu HM, 2020. Field control effect of entomopathogenic nematode against *Phyllotreta striolata* on Chinese cabbage. *China Plant Protection*, 40(11): 48–51. [杨凤丽, 曹婷婷, 张斌, 吴慧明, 2020. 昆虫病原线虫对小白菜黄曲条跳甲的田间防效. 中国植保导刊, 40(11): 48–51.]
- Yang G, Dowling AJ, Gerike U, French-Constant RH, Waterfield NR, 2006. *Photorhabdus* virulence cassettes confer injectable insecticidal activity against the wax moth. *Journal of Bacteriology*, 188(6): 2254–2261.
- Yang HW, Liu Z, Jian H, Yang XF, Chen SB, Yuan JJ, 2007. Method for culturing entomopathogenic nematodes using enzymatically hydrolyzed medium. Chinese Invention Patent, ZL200410096044.5. 2007–07–11. [杨怀文, 刘峥, 简恒, 杨秀芬, 陈松笔, 袁京京, 2007. 酶解培养基培养昆虫病原线虫的方法. 中国发明专利, ZL200410096044.5. 2007–07–11.]
- Yang HW, Zhang SG, Zhang GY, 1991. Mass production of entomopathogenic nematodes using duck intestine medium. *Bulletin of Biological Control* 7(2): 87. [杨怀文, 张善稿, 张刚应, 1991. 用鸭肠培养基大量生产昆虫病原线虫. 生物防治通报, 7(2): 87.]
- Yang J, Wang QY, Song P, Nangong ZY, Cui L, Kong FF, Feng SS, 2008. Purification and characterization of the haemocoel insecticide Tp40 from *Xenorhabdus nematophila*. *Journal of Microbiology*, 48(5): 677–683. [杨君, 王勤英, 宋萍, 南宫自艳, 崔龙, 孔繁芳, 冯姗姗, 2008. 嗜线虫致病杆菌血腔毒素 Tp40 的纯化和特性分析. 微生物学报, 48(5): 677–683.]
- Yang LK, Zhu XF, Qian XJ, Liu WH, Jiang HX, Liu CZ, 2021. Pathogenicity of five entomopathogenic nematodes from Gansu Province on *Spodoptera frugiperda*. *Pratacultural Science*, 38(10): 2069–2076. [杨淋凯, 朱小芳, 钱秀娟, 刘卫红, 姜红霞, 刘长仲, 2021. 甘肃省 5 种昆虫病原线虫对草地贪夜蛾的致病力测定. 草业科学, 38(10): 2069–2076.]
- Yang P, Li SC, 1988. Effect of temperature on the development and infectivity of entomopathogenic nematodes. *Entomological Knowledge*, 25(5): 300–302. [杨平, 李素春, 1988. 温度对昆虫病原线虫发育和感染力的影响. 昆虫知识, 25(5): 300–302.]
- Yang P, Liu J, Liu NX, Han RC, Zhang YQ, Yin CS, 2000. The application of *Steinernema* nematodes to control the *Carposina sasakii*. *Insect Natural Enemies*, 22(2): 79–82. [杨平, 刘杰, 刘南欣, 韩日畴, 张忆青, 尹纯寿, 2000. 应用斯氏线虫大面积防治桃小食心虫. 昆虫天敌, 22(2): 79–82.]
- Yang P, Xu JL, 1989. Discussion of some culture conditions affecting the output of entomogenous nematodes in mass production. *Natural Enemies of Insects*, 11(4): 187–192. [杨平, 徐洁莲, 1989. 几种培养条件对昆虫病原线虫产量和带菌率的影响. 昆虫天敌, 11(4): 187–192.]
- Yang Y, Qian X, Gao L, Hao X, Ma R, Ma Q, 2023. Effects of insulin on symbiotic bacteria of entomopathogenic nematodes. *Journal of Gansu Agricultural University*, 58(3): 133–138. [杨亚贤, 钱秀娟, 高璐, 郝欣, 马瑞鹏, 马起钰, 2023. 胰岛素对昆虫病原线虫共生菌的影响. 甘肃农业大学学报, 58(3): 133–138.]
- Yang X, Hou XQ, Sun YW, Zhang GY, Hu XF, Xie YL, Mo XT, Ding XZ, Xia LQ, Hu SB, 2019. Screening a fosmid library of *Xenorhabdus stockiae* HN_xs01 reveals SrfABC toxin that exhibits both cytotoxicity and injectable insecticidal activity. *Journal of Invertebrate Pathology*, 167: 107247.
- Yang XF, Jian H, Yang HW, Liu Z, Yuan JJ, 2004. Using entomopathogenic nematodes for control of chive maggot, *Bradysia odoriphaga* Yang et Zhang. *Journal of Plant Protection*, 31(1): 33–37. [杨秀芬, 简恒, 杨怀文, 刘峥, 袁京京, 2004. 用昆虫病原线虫防治韭蛆. 植物保护学报, 31(1): 33–37.]
- Yang XF, Yang HW, 1997. Effect of *Xenorhabdus nematophilus* strain A24 on the yield of *Steinernema carpocapsae*. *Chinese Journal of Biological Control*, 13(4): 163–165. [杨秀芬, 杨怀文, 1997. 嗜线虫杆菌接菌量对小卷蛾线虫产量的影响. 中国生物防治学报, 13(4): 163–165.]
- Yao HY, Wang LL, Wang F, Li GA, Shen JH, Lin B, 2018. An in vivo culture method for entomopathogenic nematodes. Chinese Invention Patent, ZL201511009910.7. 2018-04-06. [姚红燕, 王丽丽, 汪峰, 李国安, 湛江华, 林波, 2018. 一种昆虫病原线虫的活体繁殖方法. 中国发明专利, ZL201511009910.7. 2018-04-06.]
- Yi SW, Li XW, Chen LM, Wang JC, Huang J, Zhang J, Lv YB, 2024. Pathogenicity of entomopathogenic nematodes to red imported fire ant, *Solenopsis invicta* Buren and the control efficacy in the field. *Journal of Environmental Entomology*, 46(3): 711–717. [易松望, 李晓维, 陈利民, 王金超, 黄俊, 张娟, 吕要斌, 2024. 昆虫病原线虫对红火蚁的致病力及田间防控效果. 环境昆虫

- 学报, 46(3): 711–717.]
- Yin J, Zhu HB, Xia LQ, Ding XZ, Hoffmann T, Hoffmann M, Bian XY, Müller R, Fu J, Francis Stewart A, Zhang YM, 2015. A new recombineering system for *Photorhabdus* and *Xenorhabdus*. *Nucleic Acids Research*, 43(6): e36.
- You J, Liang SZ, Cao L, Liu XL, Han RC, 2006. Nutritive significance of crystalline inclusion proteins of *Photorhabdus luminescens* in *Steinernema* nematodes. *FEMS Microbiology Ecology*, 55(2): 178–185.
- Yu HB, Ma J, Wang RY, Geng YL, Chen SL, 2012. Virulence of entomopathogenic nematodes against sweet potato weevils. *Chinese Journal of Biological Control*, 28(4): 514–520. [于海滨, 马娟, 王容燕, 耿亚玲, 陈书龙, 2012. 昆虫病原线虫对甘薯蚜象的致病力测定. 中国生物防治学报, 28(4): 514–520.]
- Zeng WH, Chen T, Chen Y, Yan X, Wu WJ, Zhang SJ, Li ZQ, 2023. α -Terpineol affects social immunity, increasing the pathogenicity of entomopathogenic nematodes to subterranean termites (Isoptera). *Pesticide Biochemistry and Physiology*, 196: 105621.
- Zhai J, Cui LL, Wang X, Zhao LW, Zhang B, Xu LY, 2020. Effect of combination of insect-pathogenic nematodes and insecticides on *Phyllotreta striolata*. *Zhejiang Agricultural Sciences*, 61(3): 426–427. [翟婧, 崔丽利, 王笑, 赵丽稳, 张斌, 许燎原, 2020. 昆虫病原线虫与药剂组合对叶菜黄曲条跳甲的防治效果. 浙江农业科学, 61(3): 426–427.]
- Zhan F, Tian CL, Li H, Yang R, Bao H, Zhang S, Zhang X, Shi Y, Tomalak M, Půža V, Guo W, 2025. *Steinernema tarimense* n. sp. (Rhabditida: Steinernematidae), a new entomopathogenic nematode from Tarim basin, Xinjiang, China. *Journal of Helminthology*, 99: e80.
- Zhan FQ, Hou M, Yang R, Wang N, Bao HF, Hou XQ, Cui WD, Long XQ, 2020. Research on *Steinernema litorale* from *Populus euphratica* forest in Xinjiang by identification and biological characteristics. *Xinjiang Agricultural Sciences*, 57(3): 536–544. [詹发强, 侯敏, 杨蓉, 王宁, 包慧芳, 侯新强, 崔卫东, 龙宣杞, 2020. 1 株新疆胡杨林昆虫病原线虫鉴定及生物学特性. 新疆农业科学, 57(3): 536–544.]
- Zhan LL, Li CJ, Xu YL, 2008. Laboratory screening technology for *in vivo* mass production of *Steinernema carpocapsae*. *System Sciences and Comprehensive Studies in Agriculture*, 24(4): 424–426, 431. [站丽莉, 李春杰, 许艳丽, 2008. 斯氏线虫离体繁殖技术研究. 农业系统科学与综合研究, 24(4): 424–426, 431.]
- Zhang AB, 2015. Screening and evaluation of entomopathogenic nematodes and development of formulation. Master dissertation. Baoding: Hebei Agricultural University. [张安邦, 2015. 昆虫病原线虫的筛选、评价和剂型的研制. 硕士学位论文. 保定: 河北农业大学.]
- Zhang AB, Nangong ZY, Song P, Wang QY, 2015. Efficacy of entomopathogenic nematodes against *Agrotis segetum* larvae. *Journal of Environmental Entomology*, 37(3): 591–597. [张安邦, 南宫自艳, 宋萍, 王勤英, 2015. 昆虫病原线虫对黄地老虎致病力的研究. 环境昆虫学报, 37(3): 591–597.]
- Zhang C, Chen HN, Hüttel S, Hu SB, Zhang WY, Ding XZ, Yin J, Yin YL, Müller R, Xia LQ, Zhang YM, Tu Q, 2022. A novel tumor-targeting strain of *Xenorhabdus stockiae* exhibits potent biological activities. *Frontiers in Bioengineering and Biotechnology*, 10: 984197.
- Zhang CL, Wang ZM, Wang SY, Osman I, Li KB, Zhang S, Cao YZ, Wang SS, Yin J, 2024. Virulence of entomopathogenic nematodes to *Holotrichia parallela* larvae. *Chinese Journal of Biological Control*, 40(5): 1009–1014. [张彩铃, 王志敏, 王思雨, Ibrahim Osman, 李克斌, 张帅, 曹雅忠, 王森山, 尹姣, 2024. 昆虫病原线虫对暗黑鳃金龟的致病力. 中国生物防治学报, 40(5): 1009–1014.]
- Zhang HY, 2016. Screening of fine strains of entomopathogenic nematodes and analysis of its biological control. Master dissertation. Harbin: Northeast Agricultural University. [张红玉, 2016. 昆虫病原线虫优良品系的筛选及生物学特性分析. 硕士学位论文. 哈尔滨: 东北农业大学.]
- Zhang JH, Tian CL, Jiang WM, Ding Y, Zhou JC, Li MH, 2024. Advance on the effect factors of entomopathogenic nematode in control of *Cydia pomonella* L.. *Plant Quarantine*, 38(1): 13–17. [张金花, 田成丽, 姜纬明, 丁岩, 周佳春, 李茂海, 2024. 昆虫病原线虫防控苹果蠹蛾的影响因素. 植物检疫, 38(1): 13–17.]
- Zhang PB, Xu CX, Qiu XH, Han RC, 2010. Insecticidal activity and virulence of the transgenic *Enterobacter cloacae* with the insecticidal *tcdA1B1* genes against four agricultural pests. *Journal of Environmental Entomology*, 32(1): 47–53. [张鹏波, 徐彩霞, 丘雪红, 韩日畴, 2010. 转 *tcdA1B1* 杀虫基因的阴沟肠杆菌对 4 种农业害虫的杀虫活性与毒力. 环境昆虫学报, 32(1): 47–53.]
- Zhang SJ, Liu Q, Han YF, Han JH, Yan ZQ, Wang YH, Zhang X, 2019a. Nematophin, an antimicrobial dipeptide compound from *Xenorhabdus nematophila* YL001 as a potent biopesticide for *Rhizoctonia solani* control. *Frontiers in Microbiology*, 10: 1765.
- Zhang SJ, Fang XL, Tang Q, Ge J, Wang YH, Zhang X, 2019b. CpxR negatively regulates the production of xenocoumacin 1, a dihydroisocoumarin derivative produced by *Xenorhabdus nematophila*. *MicrobiologyOpen*, 8(2): e00674.

- Zhang SJ, Han YF, Wang LY, Han JH, Yan ZQ, Wang Y, Wang YH, 2024. Antifungal activity and mechanism of xenocoumacin 1, a natural product from *Xenorhabdus nematophila* against *Sclerotinia sclerotiorum*. *Journal of Fungi*, 10(3): 175.
- Zhang SJ, Qian XJ, Li CJ, Pan FJ, Xu YL, 2013. Pathogenicity of entomopathogenic nematode on *Xestia cni-grum* in soybean field. *Soybean Science*, 32(1): 63–67. [张思佳, 钱秀娟, 李春杰, 潘凤娟, 许艳丽, 2013. 昆虫病原线虫对大豆田八字地老虎幼虫致病力的研究. 大豆科学, 32(1): 63–67.]
- Zhang W, Meng J, Ning J, Qin PJ, Zhou J, Zou Z, Wang YH, Jiang H, Ahmad F, Zhao LL, Sun JH, 2017. Differential immune responses of *Monochamus alternatus* against symbiotic and entomopathogenic fungi. *Science of China: Life Sciences*, 47(9): 988–996. [张伟, 孟洁, 宁静, 秦培君, 周娇, 邹振, 王燕红, 江红, Faheem Ahmad, 赵莉莉, 孙江华, 2017. 松墨天牛对共生真菌和病原真菌的差异性免疫响应. 中国科学: 生命科学, 47(9): 988–996.]
- Zhang WW, Zhao FY, 2021. Pathogenicity of entomopathogenic nematodes to Ginkgo grubs in Enshi area. *Journal of Hunan Ecological Science*, 8(2): 40–46. [张维伟, 赵福永, 2021. 昆虫病原线虫对恩施地区银杏蛴螬的致病性研究. 湖南生态科学学报, 8(2): 40–46.]
- Zhang X, Hu XF, Li YS, Ding XZ, Yang Q, Sun YJ, Yu ZQ, Xia LQ, Hu SB, 2014. XaxAB-like binary toxin from *Photorhabdus luminescens* exhibits both insecticidal activity and cytotoxicity. *FEMS Microbiology Letters*, 350(1): 48–56.
- Zhang X, Machado RA, Doan CV, Arce CC, Hu LF, Robert CA, 2019c. Entomopathogenic nematodes increase predation success by inducing cadaver volatiles that attract healthy herbivores. *eLife*, 8: e46668.
- Zhang XB, Zhu XF, Qian XJ, 2023. Responses of *Steinernema kraussei* 0657L and *Heterorhabditis brevicaudis* 0641TY to low humidity stress. *Chinese Journal of Biological Control*, 39(2): 320–330. [张晓波, 朱小芳, 钱秀娟, 2023. 低湿胁迫下卡森氏线虫0657L和短尾异小杆线虫0641TY的避逆策略. 中国生物防治学报, 39(2): 320–330.]
- Zhang XX, 2018. Cold acclimation of *Steinernema scarabaei* and insecticidal and antibacterial activities of their symbiotic bacteria. Master dissertation. Harbin: Northeast Agricultural University. [张旭霞, 2018. 斯氏线虫低温驯化及其共生菌的杀虫抑菌活性研究. 硕士学位论文. 哈尔滨: 东北农业大学.]
- Zhang Y, Li H, Wang F, Liu C, Reddy GVP, Li H, Li ZH, Sun YC, Zhao ZH, 2023. Discovery of a new highly pathogenic toxin involved in insect sepsis. *Microbiology Spectrum*, 11(6): e01422–23.
- Zhang Y, Nangong ZY, Kong FF, Song P, Wang QY, 2013. Biological activity of *Xenorhabdus nematophila* HB310 against *Locusta migratoria* manilensis. *Journal of Pesticide Science*, 15(5): 516–522. [张园, 南宮自艳, 孔繁芳, 宋萍, 王勤英, 2013. 嗜线虫致病杆菌 HB310 菌株对东亚飞蝗的生物活性. 农药学报, 15(5): 516–522.]
- Zhang Y, Wang F, Zhao ZH, 2022. Metabonomics reveals that entomopathogenic nematodes mediate tryptophan metabolites that kill host insects. *Frontiers in Microbiology*, 13: 1042145.
- Zhang Y, Xu YL, Wu DB, Pan FJ, Xiao L, Cao YE, 2014. Management efficiency of *Bradysia odoriphaga* with entomopathogenic nematode. *Anhui Agricultural Science Bulletin*, 20(5): 70–73, 94. [张燕, 许艳丽, 武东波, 潘凤娟, 肖亮, 曹云娥, 2014. 昆虫病原线虫防治韭菜迟眼蕈蚊效果研究. 安徽农学通报, 20(5): 70–73, 94.]
- Zhang YL, Yang ZQ, Wang XY, Wu CJ, Ma SF, Lu ZG, 2012. Laboratory test on control of Japanese pine sawyer by two entomopathogenic nematode species. *Forestry Science*, 48(12): 64–70. [张彦龙, 杨忠岐, 王小艺, 吴成进, 马生福, 陆之贵, 2012. 2 种昆虫病原线虫对松褐天牛幼虫的控制力测定. 林业科学, 48(12): 64–70.]
- Zhang ZR, Cao L, Liu XL, Wang GH, Xu ZF, Han RC, 2006. Screening of insecticides as synergists for entomopathogenic nematodes *Steinernema longicaudum* X-7. *Chinese Bulletin of Entomology*, 43(1): 68–73. [张中润, 曹莉, 刘秀玲, 王果红, 许再福, 韩日畴, 2006. 昆虫病原线虫 *Steinernema longicaudum* X-7 增效药剂的筛选. 昆虫知识, 43(1): 68–73.]
- Zhang ZR, Wang JH, Huang HJ, Huang WJ, 2016. The efficacy of entomopathogenic nematodes against cashew leaf miners *Acrocercops syngamma* in the laboratory. *Journal of Environmental Entomology*, 38(3): 602–606. [张中润, 王金辉, 黄海杰, 黄伟坚, 2016. 昆虫病原线虫对腰果细蛾的致病力测定. 环境昆虫学报, 38(3): 602–606.]
- Zhang ZR, Wang JH, Huang HJ, Huang WJ, 2017. Virulence of entomopathogenic nematodes against cashew nut borer *Nephopteryx* sp. in laboratory. *Plant Protection*, 43(1): 210–213, 217. [张中润, 王金辉, 黄海杰, 黄伟坚, 2017. 昆虫病原线虫对腰果云翅斑螟的室内致病力. 植物保护, 43(1): 210–213, 217.]
- Zhao GY, 2013. Studies on application of entomopathogenic nematodes for the control of *Bradysia odoriphaga*. Master dissertation. Tai'an: Shandong Agriculture University. [赵国玉, 2013. 利用昆虫病原线虫防治韭菜迟眼蕈蚊幼虫的研究. 硕士学位论文. 泰安: 山东农业大学.]
- Zhao GY, Guo WX, 2023. Cultivation method and culture medium for *Steinernema glaseri*. Chinese Invention Patent, ZL201910760610.4. 2023-07-21. [赵国玉, 郭文秀, 2023. 昆虫病原线虫的培养方

- 法及培养基. 中国发明专利, ZL201910760610.4. 2023-07-21.]
- Zhao GY, Guo WX, Yan X, Liu YS, Chen JH, Han RC, 2013. Effects of chemical insecticides used in chive fields on the survival and infectivity of entomopathogenic nematodes. *Journal of Environmental Entomology*, 35(4): 458–465. [赵国玉, 郭文秀, 颜珣, 刘玉升, 陈镜华, 韩日畴, 2013. 韭菜田中常用化学农药对昆虫病原线虫存活及感染力的影响. 环境昆虫学报, 35(4): 458–465.]
- Zhao L, Awori RM, Kaiser M, Groß J, Opatz T, Bode HB, 2019. Structure, biosynthesis, and bioactivity of photoditritide from *Photorhabdus temperata* Meg1. *Journal of Natural Products*, 82(12): 3499–3503.
- Zhao L, Kaiser M, Bode HB, 2018. Rhabdopeptide/xenortide-like peptides from *Xenorhabdus innexi* with terminal amines showing potent antiprotozoal activity. *Organic Letters*, 20(17): 5116–5120.
- Zhao L, Le Chapelain C, Brachmann AO, Kaiser M, Groll M, Bode HB, 2021. Activation, structure, biosynthesis and bioactivity of glidobactin-like proteasome inhibitors from *Photorhabdus laumondii*. *ChemBioChem*, 22(9): 1584–1588.
- Zhao L, Vo TD, Kaiser M, Bode HB, 2020. Phototemtide A, a cyclic lipopeptide heterologously expressed from *Photorhabdus temperata* Meg1, shows selective antiprotozoal activity. *ChemBioChem*, 21(9): 1288–1292.
- Zhao LW, Fan XL, Guo HR, Ge FR, Zhai J, Ma XF, 2024. Effects of entomopathogenic nematode on controlling of *Phyllotreta striolata* in *Brassica campestris* L. ssp. *chinensis* Makino. *Journal of Changjiang Vegetables*, 2024(6): 73–75. [赵丽稳, 范雪莲, 郭焕茹, 葛芙蓉, 翟婧, 马小福, 2024. 昆虫病原线虫防治小白菜黄曲条跳甲的田间效果. 长江蔬菜, 2024(6): 73–75.]
- Zhao RH, Han RC, Qiu XH, Yan X, Cao L, Liu XL, 2008. Cloning and heterologous expression of insecticidal-protein-encoding genes from *Photorhabdus luminescens* TT01 in *Enterobacter cloacae* for termite control. *Applied and Environmental Microbiology*, 74(23): 7219–7226.
- Zhao ZY, Ru TQ, Qian ZG, Liu FL, Song XJ, 1991. Preliminary study on mass production technology of entomopathogenic nematodes. *Journal of Henan Forestry Science and Technology*, 11(3): 35–36. [赵忠懿, 茹桃勤, 钱振国, 刘凤鸾, 宋小菊, 1991. 昆虫病原线虫大量繁殖技术的初步研究. 河南林业科技, 11(3): 35–36.]
- Zhen SY, Li Y, Hou YL, Gu XH, Zhang LM, Ruan WB, Shapiro-Ilan D, 2018. Enhanced entomopathogenic nematode yield and fitness via addition of pulverized insect powder to solid media. *Journal of Nematology*, 50(4): 495–506.
- Zheng HZ, Yang KL, Qin YN, Guo WJ, Wen JB, 2021. Impacts of three chemical pesticides on infectivity of entomopathogenic nematode against *Eucryptorrhynchus brandti*. *Journal of Environmental Entomology*, 43(5): 1312–1320. [郑惠中, 杨开朗, 覃艳妮, 郭文娟, 温俊宝, 2021. 三种杀虫剂对昆虫病原线虫侵染臭椿沟眶象幼虫能力的影响. 环境昆虫学报, 43(5): 1312–1320.]
- Zheng HZ, 2021. Control test of entomopathogenic nematodes on *Eucryptorrhynchus brandti* (Harold) and *Eucryptorrhynchus scrobiculatus* Motschulsky. Master dissertation. Beijing: Beijing Forestry University. [郑惠中, 2021. 昆虫病原线虫对臭椿沟眶象和沟眶象的防治效果初试. 硕士学位论文. 北京: 北京林业大学.]
- Zhong BZ, Lü CJ, Li CX, Yan W, Huang SC, Qin WQ, 2020. Determination of lethal ability of *Heterorhabditis bacteriophora* H06 strain against the larvae of red palm weevil *Rhynchophorus ferrugineus* in laboratory. *Chinese Journal of Tropical Crops*, 41(11): 2292–2296. [钟宝珠, 吕朝军, 李朝绪, 阎伟, 黄山春, 覃伟权, 2020. 嗜菌异小杆线虫 H06 品系对红棕象甲幼虫的室内致死能力测定. 热带作物学报, 41(11): 2292–2296.]
- Zhou QQ, Bräuer A, Adihou H, Schmalhofer M, Saura P, Grammbitter GLC, Kaila VRI, Groll M, Bode HB, 2019. Molecular mechanism of polyketide shortening in anthraquinone biosynthesis of *Photorhabdus luminescens*. *Chemical Science*, 10(25): 6341–6349.
- Zhu MJ, Han RC, Wu ZQ, Liang SZ, Li QJ, Tan ZL, Li LY, Zhao RP, Cao L, Liu XL, Liu NX, 1999. The elementary research on the liquid culture of entomopathogenic nematode *Steinernema* in bioreactors. *Journal of South China University of Technology (Natural Science)*, 27(3): 83–87. [朱明军, 韩日畴, 吴振强, 梁世中, 李秋剑, 谭字榴, 李丽英, 赵汝鹏, 曹莉, 刘秀玲, 刘南欣, 1999. 斯氏线虫生物反应器液体培养的初步研究. 华南理工大学学报(自然科学版), 27(3): 83–87.]
- Zhu MJ, Tan ZL, Wu ZQ, Liang SZ, Han RC, Li QJ, Li LY, 1998. Culture of entomopathogenic nematode in bioreactors. *Chinese Journal of Biological Control*, 14(4): 167–171. [朱明军, 谭字榴, 吴振强, 梁世中, 韩日畴, 李秋剑, 李丽英, 1998. 昆虫病原线虫的生物反应器培养. 中国生物防治学报, 14(4): 167–171.]